

Boundary behavior of quotients of Martin kernels

Kentaro Hirata

Department of Mathematics, Hokkaido University,
Sapporo 060-0810, Japan *
e-mail: hirata@math.sci.hokudai.ac.jp

Abstract

Given two intersecting domains, we investigate the boundary behavior of the quotient of Martin kernels of each domain. To this end, we give a characterization of minimal thinness for a difference of two subdomains in terms of Martin kernels of each domain. As a consequence of our main theorem (Theorem 2.1), we obtain the boundary growth of the Martin kernel of a Lipschitz domain, which corresponds to earlier results for the boundary decay of the Green function for a Lipschitz domain investigated by Burdzy, Carroll and Gardiner.

Keywords: boundary behavior, Martin kernel, minimal thinness

Mathematics Subject Classifications (2000): 31B05, 31B25, 31C35

1 Introduction

One of the aims of this paper is to examine the boundary growth of the Martin kernel of a Lipschitz domain. This is motivated by earlier work due to Burdzy, Carroll and Gardiner. We write $\mathbf{0}$ for the origin of $\mathbb{R}^n$ ($n \geq 2$) to distinguish this from $0 \in \mathbb{R}$, and write $x = (x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$ and $e = (\mathbf{0}', 1)$. Suppose that $\phi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$ satisfies $\phi(\mathbf{0}') = 0$ and the Lipschitz property: there is a positive constant L such that

$$|\phi(x') - \phi(y')| \leq L|x' - y'| \quad \text{for } x', y' \in \mathbb{R}^{n-1}.$$

We put $\Omega_\phi = \{(x', x_n) : x_n > \phi(x')\}$ and set

$$I^+ = \int_{\{|x'| < 1\}} \frac{\max\{\phi(x'), 0\}}{|x'|^n} dx', \quad (1.1)$$

$$I^- = \int_{\{|x'| < 1\}} \frac{\max\{-\phi(x'), 0\}}{|x'|^n} dx'. \quad (1.2)$$

In [2, 3], Burdzy obtained a result on the angular derivative problem for analytic functions in a Lipschitz domain. The key step was to establish a relationship between the

*Current address: Faculty of Education and Human Studies, Akita University, Akita 010-8502, Japan
e-mail: hirata@math.akita-u.ac.jp

convergence of the above integrals and the boundary behavior of the Green function G_{Ω_ϕ} for Ω_ϕ . Burdzy's approach was based on probabilistic methods and the minimal fine topology. Analytic proofs were given by Carroll [4, 5] and Gardiner [7]. A proof based on the extremal length may be found in Sastry [12].

Theorem A. *Suppose that I^+ and I^- are as in (1.1) and (1.2). The following statements hold.*

(i) *If $I^+ < \infty$ and $I^- = \infty$, then*

$$\lim_{t \rightarrow 0+} \frac{G_{\Omega_\phi}(te, e)}{t} = \infty.$$

(ii) *If $I^+ = \infty$ and $I^- < \infty$, then*

$$\lim_{t \rightarrow 0+} \frac{G_{\Omega_\phi}(te, e)}{t} = 0.$$

(iii) *If $I^+ < \infty$ and $I^- < \infty$, then the limit of $G_{\Omega_\phi}(te, e)/t$, as $t \rightarrow 0+$, exists and*

$$0 < \lim_{t \rightarrow 0+} \frac{G_{\Omega_\phi}(te, e)}{t} < \infty.$$

In view of the boundary Harnack principle, Theorem A shows *the rate of boundary decay* of positive harmonic functions on Ω_ϕ vanishing continuously on a part of the boundary of Ω_ϕ near the origin. We are now interested in a relationship between the convergence of the integrals I^+, I^- and *the rate of boundary growth* of positive harmonic functions on Ω_ϕ with singularity at the origin. In view of the Fatou-Naïm-Doob theorem (see [1, Theorem 9.4.6]), it is enough to investigate it for the Martin kernel of Ω_ϕ with pole at the origin. See the first paragraph of Section 2 for the definition of the Martin kernel.

Theorem 1.1. *Suppose that I^+ and I^- are as in (1.1) and (1.2). The following statements hold.*

(i) *If $I^+ < \infty$ and $I^- = \infty$, then*

$$\lim_{t \rightarrow 0+} t^{n-1} K_{\Omega_\phi}(te, \mathbf{0}) = 0.$$

(ii) *If $I^+ = \infty$ and $I^- < \infty$, then*

$$\lim_{t \rightarrow 0+} t^{n-1} K_{\Omega_\phi}(te, \mathbf{0}) = \infty.$$

(iii) *If $I^+ < \infty$ and $I^- < \infty$, then the limit of $t^{n-1} K_{\Omega_\phi}(te, \mathbf{0})$, as $t \rightarrow 0+$, exists and*

$$0 < \lim_{t \rightarrow 0+} t^{n-1} K_{\Omega_\phi}(te, \mathbf{0}) < \infty.$$

When $I^+ = \infty$ and $I^- = \infty$, the limit of $t^{n-1}K_{\Omega_\phi}(te, \mathbf{0})$ may take the following values: 0, any positive and finite value, or ∞ .

Example 1.2. To simplify the notation, we write $\mathbb{R}_{1+}^{n-1} = \{x' \in \mathbb{R}^{n-1} : x_1 \geq 0\}$ and $\mathbb{R}_{1-}^{n-1} = \{x' \in \mathbb{R}^{n-1} : x_1 \leq 0\}$ in this example.

(i) If $\phi(x')$ is equal to $x_1/2$ on $\mathbb{R}_{1+}^{n-1}$ and x_1 on $\mathbb{R}_{1-}^{n-1}$, then

$$\lim_{t \rightarrow 0+} t^{n-1}K_{\Omega_\phi}(te, \mathbf{0}) = 0.$$

(ii) If $\phi(x')$ is equal to x_1 on $\mathbb{R}_{1+}^{n-1}$ and x_1 on $\mathbb{R}_{1-}^{n-1}$, then the limit of $t^{n-1}K_{\Omega_\phi}(te, \mathbf{0})$, as $t \rightarrow 0+$, exists and

$$0 < \lim_{t \rightarrow 0+} t^{n-1}K_{\Omega_\phi}(te, \mathbf{0}) < \infty.$$

(iii) If $\phi(x')$ is equal to x_1 on $\mathbb{R}_{1+}^{n-1}$ and $x_1/2$ on $\mathbb{R}_{1-}^{n-1}$, then

$$\lim_{t \rightarrow 0+} t^{n-1}K_{\Omega_\phi}(te, \mathbf{0}) = \infty.$$

It is easy to check that $I^+ = \infty$ and $I^- = \infty$. The value of the limit in each case follows from [9, Theorems 1 and 2].

Let $\mathbb{R}_+^n = \{(x', x_n) : x_n > 0\}$. As will be explained in Section 5, the convergence of the integrals I^+ and I^- is connected with minimal thinness of the sets $\mathbb{R}_+^n \setminus \Omega_\phi$ and $\Omega_\phi \setminus \mathbb{R}_+^n$ (see Section 2 for the definition of minimal thinness). Since $K_{\mathbb{R}_+^n}(te, \mathbf{0}) = t^{1-n}$, Theorem 1.1 may be interpreted as the relationship between minimal thinness of the sets $\mathbb{R}_+^n \setminus \Omega_\phi$, $\Omega_\phi \setminus \mathbb{R}_+^n$ and the boundary behavior of the quotient of Martin kernels of Ω_ϕ and $\mathbb{R}_+^n$. So, given two intersecting domains Φ and Ψ , it is valuable for us to investigate a relationship between minimal thinness of the differences $\Phi \setminus \Psi$, $\Psi \setminus \Phi$ and the boundary behavior of the quotient of Martin kernels of Φ and Ψ (Theorem 2.1).

2 Statement for general domains

Let Ω be a Greenian domain in $\mathbb{R}^n$ with $n \geq 2$. Here a Greenian domain means a domain possessing the Green function G_Ω for the Laplace operator. Let x_0 be a reference point in Ω . The Martin kernel of Ω is defined for $(x, y) \in (\Omega \times \Omega) \setminus \{(x_0, x_0)\}$ by

$$K_\Omega(x, y) = \frac{G_\Omega(x, y)}{G_\Omega(x_0, y)}.$$

Now, let $\{y_j\}$ be a sequence in Ω with no limit point in Ω . We observe by the Harnack inequality that if j_0 is sufficiently large, then $\{K_\Omega(\cdot, y_j)\}_{j \geq j_0}$ is a uniformly bounded sequence of positive harmonic functions on a given relatively compact open subset of Ω . Therefore the Harnack principle shows that there exists a subsequence $\{K_\Omega(\cdot, y_{j_k})\}$ converging to a positive harmonic function on Ω . The Martin boundary $\Delta(\Omega)$ of Ω is

defined as an ideal boundary consisting of all positive harmonic functions on Ω that can be obtained as the limit of $\{K_\Omega(\cdot, y_j)\}$ for some sequence $\{y_j\}$ in Ω with no limit point in Ω . The set $\Omega \cup \Delta(\Omega)$ (equipped with a suitable metric) is called a Martin compactification of Ω (see [1, Section 8.1] for details). Henceforth, we write $K_\Omega(\cdot, \xi)$ for the positive harmonic function on Ω corresponding to $\xi \in \Delta(\Omega)$. We say that $\xi \in \Delta(\Omega)$ is *minimal* if every positive harmonic function on Ω less than or equal to the corresponding Martin kernel $K_\Omega(\cdot, \xi)$ coincides with a constant multiple of $K_\Omega(\cdot, \xi)$. By $\Delta_1(\Omega)$, we denote the collection of all minimal Martin boundary points in $\Delta(\Omega)$.

The notion of minimal thinness was introduced by Naïm [11], using a regularized reduced function. Let u be a positive superharmonic function on Ω and let E be a subset of Ω . A reduced function of u relative to E on Ω is defined by

$$\Omega R_u^E(x) = \inf\{v(x)\},$$

where the infimum is taken over all positive superharmonic functions v on Ω such that $v \geq u$ on E . By $\Omega \widehat{R}_u^E$, we denote the lower semicontinuous regularization of ΩR_u^E . Observe that $\Omega \widehat{R}_u^E$ is nonnegative superharmonic on Ω and $\Omega \widehat{R}_u^E \leq u$. Let $\xi \in \Delta_1(\Delta)$. A set E is said to be *minimally thin* at ξ with respect to Ω if

$$\Omega \widehat{R}_{K_\Omega(\cdot, \xi)}^E(z) < K_\Omega(z, \xi) \quad \text{for some } z \in \Omega.$$

Minimal thinness enables us to equip the Martin compactification of Ω with the minimal fine topology. This is the collection of subsets W of the Martin compactification of Ω satisfying

- (i) $\Omega \setminus W$ is (ordinary) thin at every point of $W \cap \Omega$,
- (ii) $\Omega \setminus W$ is minimally thin at every point of $W \cap \Delta_1(\Omega)$.

See [1, Chapter 7] for details on ordinary thinness. Let U be a minimal fine neighborhood of $\xi \in \Delta_1(\Omega)$. We say that a function f on U has *minimal fine limit* l at ξ with respect to Ω if there is a subset E of Ω , minimally thin at ξ with respect to Ω , such that $f(x) \rightarrow l$ as $x \rightarrow \xi$ along $U \setminus E$, and then we write

$$\text{mf-}\lim_{\Omega} f(x) = l.$$

We note from the definition that a function is not necessarily defined on the whole of a domain when we consider the minimal fine limit.

The following theorem is our main result.

Theorem 2.1. *Let Φ and Ψ be Greenian domains in $\mathbb{R}^n$ such that $\Phi \cap \Psi$ is a non-empty domain. Suppose that $\xi \in \Delta_1(\Phi)$ is in the closure of $\Phi \cap \Psi$ in the Martin compactification of Φ , and that $\zeta \in \Delta_1(\Psi)$ is in the closure of $\Phi \cap \Psi$ in the Martin compactification of Ψ . If $\Phi \setminus \Psi$ is minimally thin at ξ with respect to Φ , then $K_\Psi(\cdot, \zeta)/K_\Phi(\cdot, \xi)$ has a finite minimal fine limit at ξ with respect to Φ . Furthermore, the following statements hold.*

- (i) *If $\Psi \setminus \Phi$ is not minimally thin at ζ with respect to Ψ , then*

$$\text{mf-}\lim_{\Phi} \frac{K_\Psi(x, \zeta)}{K_\Phi(x, \xi)} = 0.$$

(ii) If $\Psi \setminus \Phi$ is minimally thin at ζ with respect to Ψ , where ζ is the point such that

$$K_\Psi(\cdot, \zeta) - {}^\Psi R_{K_\Psi(\cdot, \zeta)}^{\Psi \setminus \Phi} = \alpha (K_\Phi(\cdot, \xi) - {}^\Phi R_{K_\Phi(\cdot, \xi)}^{\Phi \setminus \Psi}) \quad \text{on } \Phi \cap \Psi \quad (2.1)$$

for some positive constant α , then

$$0 < \text{mf-}\lim_{\Phi} \frac{K_\Psi(x, \zeta)}{K_\Phi(x, \xi)} < \infty.$$

(iii) If $\Psi \setminus \Phi$ is minimally thin at ζ with respect to Ψ , where ζ is a point such that (2.1) is not satisfied, then

$$\text{mf-}\lim_{\Phi} \frac{K_\Psi(x, \zeta)}{K_\Phi(x, \xi)} = 0.$$

We explain condition (2.1) using the following example.

Example 2.2. Let

$$\begin{aligned} \Phi &= \{(x_1, x_2) \in \mathbb{R}^2 : -2 < x_1 < 1, 0 < x_2 < 2\}, \\ \Psi &= \{(x_1, x_2) \in \mathbb{R}^2 : -1 < x_1 < 2, 0 < x_2 < 2\} \setminus E, \end{aligned}$$

where

$$E = \{(x_1, x_2) \in \mathbb{R}^2 : |x_1|^2 + |x_2 - 1/2|^2 = 1/4, x_1 > 0\}.$$

See Figure 1. Observe that the Martin boundary of Φ is homeomorphic to its Euclidean boundary and all points are minimal, and that there are two minimal Martin boundary points, say ζ_1, ζ_2 , over $(0, 0)$ with respect to Ψ . Suppose that $K_\Psi(\cdot, \zeta_1)$ is the limit of $K_\Psi(\cdot, y)$ as $y \rightarrow (0, 0)$ along $\{(0, x_2) : 0 < x_2 < 1\}$, and that $K_\Psi(\cdot, \zeta_2)$ is the limit of $K_\Psi(\cdot, y)$ as $y \rightarrow (0, 0)$ along $\{(x_1, x_2) : |x_1|^2 + |x_2 - 1|^2 = 1, x_1 > 0\}$. Also, let $\zeta_3 = (0, 2)$. Then $\Phi \setminus \Psi$ is minimally thin at $(0, 0)$ with respect to Φ , and $\Psi \setminus \Phi$ is minimally thin at ζ_j , $j = 1, 2, 3$, with respect to Ψ . It is not difficult to see that ζ_1 satisfies condition (2.1), but ζ_2 and ζ_3 do not.

For Lipschitz domains Φ and Ψ , Theorem 2.1 can be restated as the corollary below. We note from [8] that each Euclidean boundary point of a Lipschitz domain corresponds in a natural way to a unique Martin boundary point and it is minimal. So, Martin boundary points and Euclidean boundary points may be identified in this setting. Let Ω be a Lipschitz domain in $\mathbb{R}^n$ and let $c > 1$. We define a non-tangential region at $y \in \partial\Omega$ (the Euclidean boundary of Ω) by

$$\Gamma_c(y) = \{x \in \Omega : |x - y| < c \text{dist}(x, \partial\Omega)\}.$$

Note that this region is non-empty once c is sufficiently large. We say that a function f on Ω has non-tangential limit l at y if, for each c sufficiently large, $f(x)$ has limit l as $x \rightarrow y$ along $\Gamma_c(y)$. Then we write

$$\text{nt-}\lim_{\Omega} f(x) = l.$$




Figure 1: Condition (2.1)

Corollary 2.3. *Suppose that Φ and Ψ are Lipschitz domains in $\mathbb{R}^n$ such that $\Phi \cap \Psi$ is also a Lipschitz domain. Let $y \in \partial\Phi \cap \partial\Psi$, and suppose that $\Phi \setminus \Psi$ is minimally thin at y with respect to Φ . The following statements hold.*

- (i) *If $\Psi \setminus \Phi$ is not minimally thin at y with respect to Ψ , then*

$$\lim_{\substack{\text{nt} \\ \Phi \cap \Psi}} \lim_{x \rightarrow y} \frac{K_{\Psi}(x, y)}{K_{\Phi}(x, y)} = 0.$$

- (ii) *If $\Psi \setminus \Phi$ is minimally thin at y with respect to Ψ , then the non-tangential limit of $K_{\Psi}(\cdot, y)/K_{\Phi}(\cdot, y)$ at y with respect to $\Phi \cap \Psi$ exists and*

$$0 < \lim_{\substack{\text{nt} \\ \Phi \cap \Psi}} \lim_{x \rightarrow y} \frac{K_{\Psi}(x, y)}{K_{\Phi}(x, y)} < \infty.$$

Remark 2.4. If $\Phi \setminus \Psi$ is *not* minimally thin at y with respect to Φ and $\Psi \setminus \Phi$ is *not* minimally thin at y with respect to Ψ , then the non-tangential limit of $K_{\Psi}(\cdot, y)/K_{\Phi}(\cdot, y)$ may take the following values: 0, any positive and finite value, or ∞ (see Example 1.2).

3 Characterization of minimal thinness for a difference of two subdomains

Naïm [11, Théorème 11] gave a characterization of minimal thinness for a difference of two subdomains in terms of Green functions for each domain, which played an important role in the proof of Theorem A. In order to prove Theorem 2.1, we need a new characterization of minimal thinness for a difference.

Lemma 3.1. *Let Ω be a Greenian domain in $\mathbb{R}^n$ and let D be a subdomain of Ω . Suppose that $\xi \in \Delta_1(\Omega)$ is in the closure of D in the Martin compactification of Ω . The following statements are equivalent:*

(i) $\Omega \setminus D$ is minimally thin at ξ with respect to Ω ;

(ii) there exists $\eta \in \Delta_1(D)$ such that

$$\text{mf-}\lim_{D \ni x \rightarrow \eta} \frac{K_\Omega(x, \xi)}{K_D(x, \eta)} > 0. \quad (3.1)$$

Furthermore, the point $\eta \in \Delta_1(D)$ in (ii) is uniquely determined and the corresponding Martin kernel is represented as

$$K_D(\cdot, \eta) = \alpha(K_\Omega(\cdot, \xi) - {}^\Omega R_{K_\Omega(\cdot, \xi)}^{\Omega \setminus D}) \quad \text{on } D$$

for some positive constant α .

Remark 3.2. We note that the minimal fine limit in (3.1) always exists and that it satisfies

$$\text{mf-}\lim_{D \ni x \rightarrow \eta} \frac{K_\Omega(x, \xi)}{K_D(x, \eta)} = \mu_{K_\Omega(\cdot, \xi)}^D(\{\eta\}) = \inf_{x \in D} \frac{K_\Omega(x, \xi)}{K_D(x, \eta)} = \liminf_{x \rightarrow \eta} \frac{K_\Omega(x, \xi)}{K_D(x, \eta)} < \infty, \quad (3.2)$$

where $\mu_{K_\Omega(\cdot, \xi)}^D$ is the measure on $\Delta(D)$ associated with $K_\Omega(\cdot, \xi)$ in the Martin representation. See [1, Theorems 9.2.6 and 9.3.3]. Thus minimal thinness of $\Omega \setminus D$ can be also characterized in terms of any of the quantities in (3.2) instead of the minimal fine limit.

For the proof of Lemma 3.1, we need the following lemmas. Lemma 3.3 can be deduced from [1, Theorems 9.2.6 and 9.3.3]. Lemma 3.4 is due to Naïm [11, Théorème 15] (cf. [1, Theorem 9.5.5]).

Lemma 3.3. *Let E be a subset of a Greenian domain Ω in $\mathbb{R}^n$ and let $\xi \in \Delta_1(\Omega)$. The following statements are equivalent:*

(i) E is minimally thin at ξ with respect to Ω ;

(ii) there exists a positive superharmonic function u on Ω such that

$$\inf_{x \in \Omega} \frac{u(x)}{K_\Omega(x, \xi)} < \inf_{x \in E} \frac{u(x)}{K_\Omega(x, \xi)}.$$

Lemma 3.4. *Let Ω be a Greenian domain in $\mathbb{R}^n$ and let D be a subdomain of Ω . Suppose that $\xi \in \Delta_1(\Omega)$ is in the closure of D in the Martin compactification of Ω . Assume that $\Omega \setminus D$ is minimally thin at ξ with respect to Ω , and let $\eta \in \Delta_1(D)$ be the point such that*

$$K_D(\cdot, \eta) = \alpha(K_\Omega(\cdot, \xi) - {}^\Omega R_{K_\Omega(\cdot, \xi)}^{\Omega \setminus D}) \quad \text{on } D$$

for some positive constant α . The following statements for a subset E of D are equivalent:

(i) E is minimally thin at η with respect to D ;

(ii) E is minimally thin at ξ with respect to Ω .

We say that a property holds quasi-everywhere if it holds apart from a polar set. The following lemma is elementary. For the convenience of the reader, we give a proof.

Lemma 3.5. *Let D be a Greenian domain in $\mathbb{R}^n$ and let $\zeta \in \Delta_1(D)$. Then $K_D(\cdot, \zeta)$ vanishes quasi-everywhere on ∂D .*

Proof. Let V be a Martin topology (closed) neighborhood of ζ with respect to D . Then $V \cap D$ is not minimally thin at ζ with respect to D . Therefore, from [1, Theorem 6.9.1], we have

$$K_D(x, \zeta) = {}^D R_{K_D(\cdot, \zeta)}^{V \cap D}(x) = H_{K_D(\cdot, \zeta) \chi_{\partial V \cap D}}^{D \setminus V}(x) \quad \text{for } x \in D \setminus V,$$

where $H_{K_D(\cdot, \zeta) \chi_{\partial V \cap D}}^{D \setminus V}$ denotes the Perron-Wiener-Brelot solution of the Dirichlet problem in $D \setminus V$ with the boundary function $K_D(\cdot, \zeta)$ on $\partial(V \cap D) \cap D$ and 0 on ∂D . Since V is arbitrary, we obtain the lemma. $\square$

Let Ω be a domain in $\mathbb{R}^n$ and let D be a subdomain of Ω . If h is a positive harmonic function on D which vanishes quasi-everywhere on $\partial D \cap \Omega$ and is bounded near each point of $\partial D \cap \Omega$, then we see from [1, Theorem 5.2.1] that h has a subharmonic extension h^* to Ω which is valued 0 quasi-everywhere on $\partial D \cap \Omega$ and everywhere on $\Omega \setminus \overline{D}$. In what follows, we use an asterisk, as in h^* , to denote such a subharmonic extension.

Let us prove Lemma 3.1.

Proof of Lemma 3.1. By [11, Théorème 12] (cf. [1, Theorem 9.5.5]), we can easily show that (i) implies (ii). In fact, $f := K_\Omega(\cdot, \xi) - {}^\Omega R_{K_\Omega(\cdot, \xi)}^{\Omega \setminus D}$ is a minimal harmonic function on D , and so there exists $\eta \in \Delta_1(D)$ such that $K_D(\cdot, \eta) = f/f(x_0)$ on D . Hence we obtain that

$$\inf_{x \in D} \frac{K_\Omega(x, \xi)}{K_D(x, \eta)} \geq f(x_0) > 0,$$

and thus (3.1) follows from (3.2).

We next show that (ii) implies (i). We may assume that $\Omega \setminus D$ is non-polar. Let $\eta \in \Delta_1(D)$ be a point such that

$$\alpha := \text{mf-}\lim_{D} \frac{K_\Omega(x, \xi)}{K_D(x, \eta)} > 0.$$

By (3.2), we have $K_D(\cdot, \eta) \leq \alpha^{-1} K_\Omega(\cdot, \xi)$ on D . This shows that $K_D(\cdot, \eta)$ is bounded near each point of $\partial D \cap \Omega$. Also, $K_D(\cdot, \eta)$ vanishes quasi-everywhere on $\partial D \cap \Omega$ by Lemma 3.5. Thus $K_D^*(\cdot, \eta)$ is well-defined as a subharmonic function on Ω and is dominated by $\alpha^{-1} K_\Omega(\cdot, \xi)$ on Ω . Let $u = \alpha^{-1} K_\Omega(\cdot, \xi) - K_D^*(\cdot, \eta)$. Then u is superharmonic on Ω . Since $\Omega \setminus D$ is non-polar, there is a point in $\Omega \setminus D$ at which u is positive. Therefore the minimum principle yields that u is positive on Ω . Also, we find that

$$\begin{aligned} \inf_{x \in \Omega} \frac{u(x)}{K_\Omega(x, \xi)} &= \alpha^{-1} - \sup_{x \in D} \frac{K_D(x, \eta)}{K_\Omega(x, \xi)} < \alpha^{-1}, \\ \inf_{x \in \Omega \setminus (D \cup F)} \frac{u(x)}{K_\Omega(x, \xi)} &= \alpha^{-1} - \sup_{x \in \Omega \setminus (D \cup F)} \frac{K_D^*(x, \eta)}{K_\Omega(x, \xi)} = \alpha^{-1}, \end{aligned}$$

where F is a polar set in $\partial D \cap \Omega$ such that $K_D^*(\cdot, \eta) > 0$ on F . Hence it follows from Lemma 3.3 that $\Omega \setminus (D \cup F)$ is minimally thin at ξ with respect to Ω , and so is $\Omega \setminus D$.

We finally show the uniqueness of $\eta \in \Delta_1(D)$. We suppose to the contrary that there exists $\zeta \in \Delta_1(D)$ such that $K_D(\cdot, \zeta) \leq \beta K_\Omega(\cdot, \xi)$ on D and $K_D(\cdot, \zeta)$ is different from $K_D(\cdot, \eta) := \gamma(K_\Omega(\cdot, \xi) - {}^\Omega R_{K_\Omega(\cdot, \xi)}^{\Omega \setminus D})$, where β and γ are some positive constants. We may assume that β is the smallest number satisfying $K_D(\cdot, \zeta) \leq \beta K_\Omega(\cdot, \xi)$ on D . Since $\xi \in \Delta_1(\Omega)$, it follows that $\beta K_\Omega(\cdot, \xi)$ is the least harmonic majorant of $K_D^*(\cdot, \zeta)$ on Ω . Let W be a Martin topology neighborhood of ζ with respect to D such that η is apart from W . Then $W \cap D$ is minimally thin at η with respect to D . Thus minimal thinness of $\Omega \setminus D$ at ξ with respect to Ω , together with Lemma 3.4, yields that $W \cap D$ is minimally thin at ξ with respect to Ω .

On the other hand, since $W \cap D$ is not minimally thin at ζ with respect to D , we have that

$$K_D(\cdot, \zeta) = {}^D \widehat{R}_{K_D(\cdot, \zeta)}^{W \cap D} \leq \beta {}^D \widehat{R}_{K_\Omega(\cdot, \xi)}^{W \cap D} \leq \beta {}^\Omega \widehat{R}_{K_\Omega(\cdot, \xi)}^{W \cap D} \quad \text{on } D.$$

Since $\beta K_\Omega(\cdot, \xi)$ is the least one among superharmonic functions u on Ω satisfying $K_D^*(\cdot, \zeta) \leq u$ on Ω , we have ${}^\Omega \widehat{R}_{K_\Omega(\cdot, \xi)}^{W \cap D} = K_\Omega(\cdot, \xi)$ on Ω , so that $W \cap D$ is not minimally thin at ξ with respect to Ω . Thus we obtain a contradiction, and hence the uniqueness of $\eta \in \Delta_1(D)$ is established. The proof of Lemma 3.1 is complete. $\square$

4 Proofs of Theorem 2.1 and Corollary 2.3

We give proofs of Theorem 2.1 and Corollary 2.3.

Proof of Theorem 2.1. In order to prove the first assertion, we assume that $\Phi \setminus (\Phi \cap \Psi)$ is minimally thin at ξ with respect to Φ . Let $\eta \in \Delta_1(\Phi \cap \Psi)$ be the point such that

$$K_{\Phi \cap \Psi}(\cdot, \eta) = \alpha(K_\Phi(\cdot, \xi) - {}^\Phi R_{K_\Phi(\cdot, \xi)}^{\Phi \setminus \Psi})$$

on $\Phi \cap \Psi$ for some positive constant α . Then, by Lemma 3.1 with $D := \Phi \cap \Psi$ and $\Omega := \Phi$ and Remark 3.2, we find that the minimal fine limit of $K_\Phi(\cdot, \xi)/K_{\Phi \cap \Psi}(\cdot, \eta)$ at η with respect to $\Phi \cap \Psi$ exists and

$$0 < \text{mf} - \lim_{\Phi \cap \Psi} \frac{K_\Phi(x, \xi)}{K_{\Phi \cap \Psi}(x, \eta)} < \infty. \quad (4.1)$$

It also follows from [1, Theorem 9.3.3] that $K_\Psi(\cdot, \zeta)/K_{\Phi \cap \Psi}(\cdot, \eta)$ has a finite minimal fine limit at η with respect to $\Phi \cap \Psi$. The minimal thinness of $\Phi \setminus (\Phi \cap \Psi)$ at ξ with respect to Φ , together with Lemma 3.4 with $D := \Phi \cap \Psi$ and $\Omega := \Phi$, allows us to conclude that $K_\Psi(\cdot, \zeta)/K_\Phi(\cdot, \xi)$ has a finite minimal fine limit at ξ with respect to Φ .

To prove (i), we assume in addition that $\Psi \setminus (\Phi \cap \Psi)$ is not minimally thin at ζ with respect to Ψ . Then Lemma 3.1 with $D := \Phi \cap \Psi$ and $\Omega := \Psi$ shows that for any $\eta \in \Delta_1(\Phi \cap \Psi)$, the minimal fine limit in (3.1) is zero. Therefore we have that

$$\text{mf} - \lim_{\Phi \cap \Psi} \frac{K_\Psi(x, \zeta)}{K_{\Phi \cap \Psi}(x, \eta)} = 0.$$

Hence (i) follows from (4.1) and Lemma 3.4 with $D := \Phi \cap \Psi$ and $\Omega := \Phi$.

To prove (ii), we assume in addition that $\Psi \setminus (\Phi \cap \Psi)$ is minimally thin at ζ with respect to Ψ , where ζ is the point in $\Delta_1(\Psi)$ such that (2.1) is satisfied. We note from (2.1) that $K_{\Phi \cap \Psi}(\cdot, \eta)$ is also written as $\beta(K_\Psi(\cdot, \zeta) - {}^\Psi R_{K_\Psi(\cdot, \zeta)}^{\Psi \setminus \Phi})$ on $\Phi \cap \Psi$ for some positive constant β . Then, by Lemma 3.1 with $D := \Phi \cap \Psi$ and $\Omega := \Psi$ and Remark 3.2, we find that the minimal fine limit of $K_\Psi(\cdot, \zeta)/K_{\Phi \cap \Psi}(\cdot, \eta)$ at η with respect to $\Phi \cap \Psi$ exists and

$$0 < \text{mf} - \lim_{\Phi \cap \Psi} \frac{K_\Psi(x, \zeta)}{K_{\Phi \cap \Psi}(x, \eta)} < \infty.$$

Therefore (ii) follows from (4.1) and Lemma 3.4 with $D := \Phi \cap \Psi$ and $\Omega := \Phi$.

To prove (iii), we assume in addition that $\Psi \setminus (\Phi \cap \Psi)$ is minimally thin at ζ with respect to Ψ , where ζ is a point in $\Delta_1(\Psi)$ such that (2.1) is not satisfied. Then the normalization $K_{\Phi \cap \Psi}(\cdot, \omega)$ of $K_\Psi(\cdot, \zeta) - {}^\Psi R_{K_\Psi(\cdot, \zeta)}^{\Psi \setminus \Phi}$ at a reference point is a minimal Martin kernel of $\Phi \cap \Psi$, but is different from $K_{\Phi \cap \Psi}(\cdot, \eta)$. We note from the uniqueness in Lemma 3.1 that for only $\omega \in \Delta_1(\Phi \cap \Psi)$, $K_\Psi(\cdot, \zeta)/K_{\Phi \cap \Psi}(\cdot, \omega)$ has a positive minimal fine limit at ω with respect to $\Phi \cap \Psi$. Therefore we have that

$$\text{mf} - \lim_{\Phi \cap \Psi} \frac{K_\Psi(x, \zeta)}{K_{\Phi \cap \Psi}(x, \eta)} = 0.$$

Hence (iii) follows from (4.1) and Lemma 3.4 with $D := \Phi \cap \Psi$ and $\Omega := \Phi$. Thus Theorem 2.1 is established. $\square$

Proof of Corollary 2.3. Let $y \in \partial\Phi \cap \partial\Psi$. We first show (i). By Theorem 2.1 (i) and Lemma 3.4, we find that $K_\Psi(\cdot, y)/K_\Phi(\cdot, y)$ has minimal fine limit 0 at y with respect to $\Phi \cap \Psi$. Since the non-tangential region $\Gamma_c(y)$ is not minimally thin at y with respect to $\Phi \cap \Psi$ (see [8, Section 5]), the existence of the minimal fine limit of $K_\Psi(\cdot, y)/K_\Phi(\cdot, y)$ with respect to $\Phi \cap \Psi$ implies the existence of the non-tangential limit with respect to $\Phi \cap \Psi$, and the both values coincide. Hence (i) follows.

We next show (ii). We observe that $K_\Phi(\cdot, y)$ and $K_\Psi(\cdot, y)$ satisfy (2.1) on $\Phi \cap \Psi$, since $K_\Phi(\cdot, y) - {}^\Phi R_{K_\Phi(\cdot, y)}^{\Phi \setminus \Psi}$ and $K_\Psi(\cdot, y) - {}^\Psi R_{K_\Psi(\cdot, y)}^{\Psi \setminus \Phi}$ are minimal harmonic functions on $\Phi \cap \Psi$ with pole at y . Therefore (ii) follows from Theorem 2.1 (ii). $\square$

5 Proof of Theorem 1.1

In order to prove Theorem 1.1, we collect lemmas on relationships between the convergence of the integrals I^+ , I^- in (1.1), (1.2) and minimal thinness of the differences $\Omega_\phi \setminus \mathbb{R}_+^n$, $\mathbb{R}_+^n \setminus \Omega_\phi$. See [7, Lemma 1 and Proof of Theorem 1] for Lemma 5.1 and [6, Theorem 4.2] for Lemma 5.2.

Lemma 5.1. *The following statements hold.*

- (i) $I^+ < \infty$ if and only if $\mathbb{R}_+^n \setminus \Omega_\phi$ is minimally thin at $\mathbf{0}$ with respect to $\mathbb{R}_+^n$.
- (ii) If $I^+ < \infty$ and $I^- = \infty$, then $\Omega_\phi \setminus \mathbb{R}_+^n$ is not minimally thin at $\mathbf{0}$ with respect to Ω_ϕ .

Lemma 5.2. *Let Ω be a Greenian domain in $\mathbb{R}^n$ containing $\mathbb{R}_+^n$. Suppose that Ω has a unique Martin boundary point at infinity and it is minimal. If $\Omega \setminus \overline{\mathbb{R}_+^n}$ is minimally thin at ∞ with respect to $\mathbb{R}_-^n := \{(x', x_n) : x_n < 0\}$, then $\Omega \setminus \mathbb{R}_+^n$ is minimally thin at ∞ with respect to Ω .*

Lemma 5.3. *If $I^- < \infty$, then $\Omega_\phi \setminus \mathbb{R}_+^n$ is minimally thin at $\mathbf{0}$ with respect to $\Omega_\phi \cup \mathbb{R}_+^n$.*

Proof. By Lemma 5.1, we see that $\Omega_\phi \setminus \overline{\mathbb{R}_+^n}$ is minimally thin at $\mathbf{0}$ with respect to $\mathbb{R}_-^n$. Since minimal thinness is invariant under the inversion with respect to the unit sphere, it follows from Lemma 5.2 that $\Omega_\phi \setminus \mathbb{R}_+^n$ is minimally thin at $\mathbf{0}$ with respect to $\Omega_\phi \cup \mathbb{R}_+^n$. $\square$

Lemma 5.4. *If $I^+ < \infty$ and $I^- < \infty$, then $\Omega_\phi \setminus \mathbb{R}_+^n$ is minimally thin at $\mathbf{0}$ with respect to Ω_ϕ .*

Proof. We note from Lemma 5.3 that $(\Omega_\phi \cup \mathbb{R}_+^n) \setminus \mathbb{R}_+^n$ is minimally thin at $\mathbf{0}$ with respect to $\Omega_\phi \cup \mathbb{R}_+^n$. Therefore we see from Lemmas 3.4 and 5.1 that $(\Omega_\phi \cup \mathbb{R}_+^n) \setminus \Omega_\phi$ is minimally thin at $\mathbf{0}$ with respect to $\Omega_\phi \cup \mathbb{R}_+^n$. Applying Lemma 3.4 again, we obtain the lemma. $\square$

Let us now prove Theorem 1.1.

Proof of Theorem 1.1. We can easily obtain (i) and (iii) from Corollary 2.3 with $\Phi := \mathbb{R}_+^n$ and $\Psi := \Omega_\phi$ and Lemmas 5.1 and 5.4. We show (ii). Since $(\Omega_\phi \cup \mathbb{R}_+^n) \setminus \mathbb{R}_+^n$ is minimally thin at $\mathbf{0}$ with respect to $\Omega_\phi \cup \mathbb{R}_+^n$ by Lemma 5.3, it follows from Lemma 3.1 with $D := \mathbb{R}_+^n$ and $\Omega := \Omega_\phi \cup \mathbb{R}_+^n$ that $K_{\Omega_\phi \cup \mathbb{R}_+^n}(\cdot, \mathbf{0})/K_{\mathbb{R}_+^n}(\cdot, \mathbf{0})$ has a positive minimal fine limit at $\mathbf{0}$ with respect to $\mathbb{R}_+^n$. Therefore $t^{n-1}K_{\Omega_\phi \cup \mathbb{R}_+^n}(te, \mathbf{0})$ has a positive limit as $t \rightarrow 0+$. Also, it follows from Lemmas 3.4 and 5.1 that $(\Omega_\phi \cup \mathbb{R}_+^n) \setminus \Omega_\phi$ is not minimally thin at $\mathbf{0}$ with respect to $\Omega_\phi \cup \mathbb{R}_+^n$. Therefore we have by Lemma 3.1 with $D := \Omega_\phi$ and $\Omega := \Omega_\phi \cup \mathbb{R}_+^n$ that $K_{\Omega_\phi \cup \mathbb{R}_+^n}(\cdot, \mathbf{0})/K_{\Omega_\phi}(\cdot, \mathbf{0})$ has minimal fine limit 0 at $\mathbf{0}$ with respect to Ω_ϕ , and so $K_{\Omega_\phi}(te, \mathbf{0})/K_{\Omega_\phi \cup \mathbb{R}_+^n}(te, \mathbf{0})$ has limit ∞ as $t \rightarrow 0+$. Thus we conclude that $t^{n-1}K_{\Omega_\phi}(te, \mathbf{0})$ has limit ∞ as $t \rightarrow 0+$. $\square$

Remark 5.5. For a non-Lipschitz function ϕ , a relationship between the convergence of the integral I^+ and minimal thinness of the difference $\Omega_\phi \setminus \overline{\mathbb{R}_+^n}$ was recently studied in [10].

Acknowledgement

The author is very grateful to the referee for helpful comments and suggestions.

Note added in the proof

Recently, we have obtained a relationship between the boundary decay rate of the Green function and the growth rate of the Martin kernel (Estimates for the products of the Green function and the Martin kernel, Nagoya Math. J., to appear).

References

- [1] D. H. Armitage and S. J. Gardiner, *Classical potential theory*, Springer-Verlag London, 2001.
- [2] K. Burdzy, *Brownian excursions and minimal thinness. Part III: Applications to the angular derivative problem*, Math. Z. **192** (1986), no. 1, 89–107.
- [3] ———, *Multidimensional Brownian excursions and potential theory*, Pitman Research Notes in Mathematics Series, **164**, 1987.
- [4] T. Carroll, *A classical proof of Burdzy's theorem on the angular derivative*, J. London Math. Soc. (2) **38** (1988), no. 3, 423–441.
- [5] ———, *Boundary behaviour of positive harmonic functions on Lipschitz domains*, Ann. Acad. Sci. Fenn. Math. **27** (2002), no. 1, 231–236.
- [6] V. Eiderman and M. Essén, *Harmonic majorization of $|x_1|$ in subsets of $\mathbf{R}^n$, $n \geq 2$* , Ann. Acad. Sci. Fenn. Math., **21** (1996), no. 1, 223–240.
- [7] S. J. Gardiner, *A short proof of Burdzy's theorem on the angular derivative*, Bull. London Math. Soc. **23** (1991), no. 6, 575–579.
- [8] R. A. Hunt and R. L. Wheeden, *Positive harmonic functions on Lipschitz domains*, Trans. Amer. Math. Soc. **147** (1970), 507–527.
- [9] Ü. Kuran, *On NTA-conical domains*, J. London Math. Soc. (2) **40** (1989), no. 3, 467–475.
- [10] T. Lundh, *Minimally thin sets below a function graph*, Complex Variables Theory Appl. **49** (2004), no. 7–9, 639–645.
- [11] L. Naïm, *Sur le rôle de la frontière de R. S. Martin dans la théorie du potentiel*, Ann. Inst. Fourier (Grenoble) **7**, 1957, 183–281.
- [12] S. Sastry, *Existence of angular derivative for a class of strip domains*, Proc. Amer. Math. Soc. **123** (1995), no. 4, 1075–1082.