

CLASSIFICATION OF EXTREMAL ELLIPTIC $K3$ SURFACES AND FUNDAMENTAL GROUPS OF OPEN $K3$ SURFACES

ICHIRO SHIMADA AND DE-QI ZHANG

ABSTRACT. We present a complete list of extremal elliptic $K3$ surfaces (Theorem 1.1). As an application, we give a sufficient condition for the topological fundamental group of complement to an ADE -configuration of smooth rational curves on a $K3$ surface to be trivial (Proposition 4.1 and Theorems 4.3).

1. INTRODUCTION

A complex elliptic $K3$ surface $f : X \rightarrow \mathbb{P}^1$ with a section O is said to be *extremal* if the Picard number $\rho(X)$ of X is 20 and the Mordell-Weil group MW_f of f is finite. The purpose of this paper is to present the complete list of all extremal elliptic $K3$ surfaces. As an application, we show that, if an ADE -configuration of smooth rational curves on a $K3$ surface satisfies a certain condition, then the topological fundamental group of the complement is trivial. (See Theorem 4.3 for the precise statement.)

Let $f : X \rightarrow \mathbb{P}^1$ be an elliptic $K3$ surface with a section O . We denote by R_f the set of all points $v \in \mathbb{P}^1$ such that $f^{-1}(v)$ is reducible. For a point $v \in R_f$, let $f^{-1}(v)^\#$ be the union of irreducible components of $f^{-1}(v)$ that are disjoint from the zero section O . It is known that the cohomology classes of irreducible components of $f^{-1}(v)^\#$ form a negative definite root lattice $S_{f,v}$ of type A_l , D_m or E_n in $H^2(X; \mathbb{Z})$. Let $\tau(S_{f,v})$ be the type of this lattice. We define Σ_f to be the formal sum of these types;

$$\Sigma_f := \sum_{v \in R_f} \tau(S_{f,v}).$$

The Néron-Severi lattice NS_X of X is defined to be $H^{1,1}(X) \cap H^2(X; \mathbb{Z})$, and the transcendental lattice T_X of X is defined to be the orthogonal complement of NS_X in $H^2(X; \mathbb{Z})$. We call the triple (Σ_f, MW_f, T_X) the *data* of the elliptic $K3$ surface $f : X \rightarrow \mathbb{P}^1$. When $f : X \rightarrow \mathbb{P}^1$ is extremal, the transcendental lattice T_X is a positive definite even lattice of rank 2.

Theorem 1.1. *There exists an extremal elliptic $K3$ surface $f : X \rightarrow \mathbb{P}^1$ with data (Σ_f, MW_f, T_X) if and only if (Σ_f, MW_f, T_X) appears in Table 2 given at the end of this paper.*

In Table 2, the transcendental lattice T_X is expressed by the coefficients of its Gram matrix

$$\begin{pmatrix} a & b \\ b & c \end{pmatrix}.$$

See Subsection 2.1 on how to recover the $K3$ surface X from T_X .

The classification of *semi-stable* extremal elliptic $K3$ surfaces has been done by Miranda and Persson[7] and complemented by Artal-Bartolo, Tokunaga and Zhang[1]. We can check that the semi-stable part of our list (No. 1- No. 112) coincides with theirs. Nishiyama[12] classified all elliptic fibrations (not necessarily extremal) on certain $K3$ surfaces. On the other hand, Ye[19] has independently classified all extremal elliptic $K3$ surfaces with no semi-stable singular fibers by different methods from ours.

Acknowledgment. The authors would like to thank Professors Shigeyuki Kondō, Ken-ichi Nishiyama and Keiji Oguiso for helpful discussions.

2. PRELIMINARIES

2.1. Transcendental lattice of singular $K3$ surfaces. Let \mathcal{Q} be the set of symmetric matrices

$$Q = \begin{pmatrix} a & b \\ b & c \end{pmatrix}$$

of integer coefficients such that a and c are even and that the corresponding quadratic forms are positive definite. The group $GL_2(\mathbb{Z})$ acts on \mathcal{Q} from right by

$$Q \mapsto {}^t g \cdot Q \cdot g,$$

where $g \in GL_2(\mathbb{Z})$. Let Q_1 and Q_2 be two matrices in \mathcal{Q} , and let L_1 and L_2 be the positive definite even lattices of rank 2 whose Gram matrices are Q_1 and Q_2 , respectively. Then L_1 and L_2 are isomorphic as lattices if and only if Q_1 and Q_2 are in the same orbit under the action of $GL_2(\mathbb{Z})$. On the other hand, each orbit in \mathcal{Q} under the action of $SL_2(\mathbb{Z})$ contains a unique matrix with coefficients satisfying

$$-a < 2b \leq a \leq c, \quad \text{with } b \geq 0 \text{ if } a = c.$$

(See, for example, Conway and Sloane[3, p. 358].) Hence each orbit in \mathcal{Q} under the action of $GL_2(\mathbb{Z})$ contains a unique matrix with coefficients satisfying

$$(2.1) \quad 0 \leq 2b \leq a \leq c.$$

In Table 2, the transcendental lattice is represented by the Gram matrix satisfying the condition(2.1).

Let X be a $K3$ surface with $\rho(X) = 20$; that is, X is a singular $K3$ surface in the terminology of Shioda and Inose[16]. The transcendental lattice T_X can be naturally oriented by means of a holomorphic two form on X (cf. [16, p.128]). Let \mathcal{S} denote the set of isomorphism classes of singular $K3$ surfaces. Using the natural orientation on the transcendental lattice, we can lift the map $\mathcal{S} \rightarrow \mathcal{Q}/GL_2(\mathbb{Z})$ given by $X \mapsto T_X$ to the map $\mathcal{S} \rightarrow \mathcal{Q}/SL_2(\mathbb{Z})$.

Proposition 2.1 (Shioda and Inose[16]). *This map $\mathcal{S} \rightarrow \mathcal{Q}/SL_2(\mathbb{Z})$ is bijective.* \square

Moreover, Shioda and Inose[16] gave us a method to construct explicitly the singular $K3$ surface corresponding to a given element of $\mathcal{Q}/SL_2(\mathbb{Z})$ by means of Kummer surfaces. The injectivity of the map $\mathcal{S} \rightarrow \mathcal{Q}/SL_2(\mathbb{Z})$ had been proved by Piateskii-Shapiro and Shafarevich[14].

Suppose that an orbit $[Q] \in \mathcal{Q}/GL_2(\mathbb{Z})$ is represented by a matrix Q satisfying (2.1). Let $\rho : \mathcal{Q}/SL_2(\mathbb{Z}) \rightarrow \mathcal{Q}/GL_2(\mathbb{Z})$ be the natural projection. Then we

have

$$|\rho^{-1}([Q])| = \begin{cases} 2 & \text{if } 0 < 2b < a < c \\ 1 & \text{otherwise.} \end{cases}$$

Therefore, if a data in Table 2 satisfies $a = c$ or $b = 0$ or $2b = a$ (resp. $0 < 2b < a < c$), then the number of the isomorphism classes of $K3$ surfaces that possess a structure of the extremal elliptic $K3$ surfaces with the given data is one (resp. two).

2.2. Roots of a negative definite even lattice. Let M be a negative definite even lattice. A vector of M is said to be a *root* of M if its norm is -2 . We denote by $\text{root}(M)$ the number of roots of M , and by M_{root} the sublattice of M generated by the roots of M . Suppose that a Gram matrix (a_{ij}) of M is given. Then $\text{root}(M)$ can be calculated by the following method. Let

$$g_r(x) = - \sum_{i,j=1}^r a_{ij} x_i x_j$$

be the positive definite quadratic form associated with the opposite lattice M^- of M , where r is the rank of M . We consider the bounded closed subset

$$E(g_r, 2) := \{x \in \mathbb{R}^r ; g_r(x) \leq 2\}$$

of \mathbb{R}^r . Then we have

$$\text{root}(M) + 1 = |E(g_r, 2) \cap \mathbb{Z}^r|,$$

where $+1$ comes from the origin. For a positive integer k less than r , we write by $p_k : \mathbb{R}^r \rightarrow \mathbb{R}^k$ the projection $(x_1, \dots, x_r) \mapsto (x_1, \dots, x_k)$. Then there exists a positive definite quadratic form g_k of variables (x_1, \dots, x_k) and a positive real number σ_k such that

$$p_k(E(g_r, 2)) = E(g_k, \sigma_k) := \{y \in \mathbb{R}^k ; g_k(y) \leq \sigma_k\}.$$

The projection $(x_1, \dots, x_{k+1}) \mapsto (x_1, \dots, x_k)$ maps $E(g_{k+1}, \sigma_{k+1})$ to $E(g_k, \sigma_k)$. Hence, if we have the list of the points of $E(g_k, \sigma_k) \cap \mathbb{Z}^k$, then it is easy to make the list of the points of $E(g_{k+1}, \sigma_{k+1}) \cap \mathbb{Z}^{k+1}$. Thus, starting from $E(g_1, \sigma_1) \cap \mathbb{Z}$, we can make the list of the points of $E(g_r, 2) \cap \mathbb{Z}^r$ by induction on k .

2.3. Root lattices of type ADE . A *root type* is, by definition, a finite formal sum Σ of A_l , D_m and E_n with non-negative integer coefficients;

$$\Sigma = \sum_{l \geq 1} a_l A_l + \sum_{m \geq 4} d_m D_m + \sum_{n=6}^8 e_n E_n.$$

We denote by $L(\Sigma)$ the negative definite root lattice corresponding to Σ . The rank of $L(\Sigma)$ is given by

$$\text{rank}(L(\Sigma)) = \sum_{l \geq 1} a_l l + \sum_{m \geq 4} d_m m + \sum_{n=6}^8 e_n n,$$

and the number of roots of $L(\Sigma)$ is given by

$$(2.2) \quad \text{root}(L(\Sigma)) = \sum_{l \geq 1} a_l (l^2 + l) + \sum_{m \geq 4} d_m (2m^2 - 2m) + 72e_6 + 126e_7 + 240e_8.$$

(See, for example, Bourbaki[2].) Because of $L(\Sigma)_{root} = L(\Sigma)$, we have

$$(2.3) \quad L(\Sigma_1) \cong L(\Sigma_2) \iff \Sigma_1 = \Sigma_2.$$

We also define $eu(\Sigma)$ by

$$eu(\Sigma) := \sum_{l \geq 1} a_l(l+1) + \sum_{m \geq 4} d_m(m+2) + \sum_{n=6}^8 e_n(n+2).$$

Lemma 2.2. *Let $f : X \rightarrow \mathbb{P}^1$ be an elliptic K3 surface. Then $eu(\Sigma_f)$ is at most 24. Moreover, if $eu(\Sigma_f) < 24$, then there exists at least one singular fiber of type I₁, II, III or IV.*

Proof. Let $e(Y)$ denote the topological euler number of a CW-complex Y . Then $e(X) = 24$ is equal to the sum of topological euler numbers of singular fibers of f . Every singular fiber has a positive topological euler number. We have defined $eu(\Sigma)$ in such a way that, if $v \in R_f$, then $eu(\tau(S_{f,v})) \leq e(f^{-1}(v))$ holds, and if $eu(\tau(S_{f,v})) < e(f^{-1}(v))$, then the type of the fiber $f^{-1}(v)$ is either III or IV. Hence $eu(\Sigma_f)$ does not exceed the sum of the topological euler numbers of reducible singular fibers, and if $eu(\Sigma_f) < 24$, then there is an irreducible singular fiber or a singular fiber of type III or IV. \square

2.4. Discriminant form and overlattices. Let L be an even lattice, L^\vee the dual of L , D_L the discriminant group L^\vee/L of L , and q_L the discriminant form on D_L . (See Nikulin[11, n. 4] for the definitions.) An overlattice of L is, by definition, an integral sublattice of the \mathbb{Q} -lattice L^\vee containing L .

Lemma 2.3 (Nikulin[11] Proposition 1.4.2). (1) *Let A be an isotropic subgroup of (D_L, q_L) . Then the pre-image $M := \phi_L^{-1}(A)$ of A by the natural projection $\phi_L : L^\vee \rightarrow D_L$ is an overlattice of L , and the discriminant form (D_M, q_M) of M is isomorphic to $(A^\perp/A, q_L|_{A^\perp/A})$, where A^\perp is the orthogonal complement of A in D_L , and $q_L|_{A^\perp/A}$ is the restriction of q_L to A^\perp/A .* (2) *The correspondence $A \mapsto M$ gives a bijection from the set of isotropic subgroups of (D_L, q_L) to the set of even overlattices of L .* \square

Lemma 2.4 (Nikulin[11] Corollary 1.6.2). *Let S and K be two even lattices. Then the following two conditions are equivalent. (i) There is an isomorphism $\gamma : D_S \xrightarrow{\sim} D_K$ of abelian groups such that $\gamma^*q_K = -q_S$. (ii) There is an even unimodular overlattice of $S \oplus K$ into which S and K are primitively embedded.* \square

2.5. Néron-Severi groups of elliptic K3 surfaces. Let $f : X \rightarrow \mathbb{P}^1$ be an elliptic K3 surface with the zero section O . In the Néron-Severi lattice NS_X of X , the cohomology classes of the zero section O and a general fiber of f generate a sublattice U_f of rank 2, which is isomorphic to the hyperbolic lattice

$$H := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Let W_f be the orthogonal complement of U_f in NS_X . Because U_f is unimodular, we have $NS_X = U_f \oplus W_f$. Because U_f is of signature $(1, 1)$ and NS_X is of signature $(1, \rho(X) - 1)$, W_f is negative definite of rank $\rho(X) - 2$. Note that W_f contains the sublattice

$$S_f := \bigoplus_{v \in R_f} S_{f,v}$$

generated by the cohomology classes of irreducible components of reducible fibers of f that are disjoint from the zero section. By definition, S_f is isomorphic to $L(\Sigma_f)$.

Lemma 2.5 (Nishiyama[12] Lemma 6.1). *The sublattice S_f of W_f coincides with $(W_f)_{\text{root}}$, and the Mordell-Weil group MW_f of f is isomorphic to W_f/S_f . In particular, $\text{root}(L(\Sigma_f))$ is equal to $\text{root}(W_f)$. \square*

Because $W_f \oplus U_f \oplus T_X$ has an even unimodular overlattice $H^2(X; \mathbb{Z})$ into which $NS_X = W_f \oplus U_f$ and T_X are primitively embedded, and because the discriminant form of NS_X is equal to the discriminant form of W_f by $D_{U_f} = (0)$, Lemma 2.4 implies the following:

Corollary 2.6. *There is an isomorphism $\gamma : D_{W_f} \xrightarrow{\sim} D_{T_X}$ of abelian groups such that $\gamma^* q_{T_X}$ coincides with $-q_{W_f}$. \square*

2.6. Existence of elliptic K3 surfaces. Let Λ be the K3 lattice $L(2E_8) \oplus H^{\oplus 3}$.

Lemma 2.7 (Kondō[5] Lemma 2.1). *Let T be a positive definite primitive sublattice of Λ with $\text{rank}(T) = 2$, and T^\perp the orthogonal complement of T in Λ . Suppose that T^\perp contains a sublattice H_T isomorphic to the hyperbolic lattice. Let M_T be the orthogonal complement of H_T in T^\perp . Then there exists an elliptic K3 surface $f : X \rightarrow \mathbb{P}^1$ such that $T_X \cong T$ and $W_f \cong M_T$.*

Proof. By the surjectivity of the period map of the moduli of K3 surfaces (cf. Todorov[17]), there exist a K3 surface X and an isomorphism $\alpha : H^2(X; \mathbb{Z}) \cong \Lambda$ of lattices such that $\alpha^{-1}(T) = T_X$. By Kondō[5, Lemma 2.1], the K3 surface X has an elliptic fibration $f : X \rightarrow \mathbb{P}^1$ with a section such that $\mathbb{Z}[F]^\perp/\mathbb{Z}[F] \cong M_T$, where $[F] \in U_f$ is the cohomology class of a fiber of f , and $\mathbb{Z}[F]^\perp$ is the orthogonal complement of $[F]$ in the Néron-Severi lattice NS_X . Because NS_X coincides with $U_f \oplus W_f$, and because $\mathbb{Z}[F]^\perp \cap U_f$ coincides with $\mathbb{Z}[F]$, we see that $\mathbb{Z}[F]^\perp/\mathbb{Z}[F]$ is isomorphic to W_f . \square

2.7. Datum of extremal elliptic K3 surfaces.

Proposition 2.8. *A triple (Σ, MW, T) consisting of a root type Σ , a finite abelian group MW and a positive definite even lattice T of rank 2 is a data of an extremal elliptic K3 surface if and only if the following hold:*

(D1) $\text{length}(MW) \leq 2$, $\text{rank}(L(\Sigma)) = 18$ and $eu(\Sigma) \leq 24$.

(D2) *There exists an overlattice M of $L(\Sigma)$ satisfying the following:*

(D2-a) $M/L(\Sigma) \cong MW$,

(D2-b) *there exists an isomorphism $\gamma : D_M \xrightarrow{\sim} D_T$ of abelian groups such that $\gamma^* q_T = -q_M$, and*

(D2-c) $\text{root}(L(\Sigma)) = \text{root}(M)$.

Proof. Suppose that there exists an extremal elliptic K3 surface $f : X \rightarrow \mathbb{P}^1$ with data equal to (Σ, MW, T) . It is obvious that Σ and MW satisfies the condition (D1). Via the isomorphism $S_f \cong L(\Sigma)$, the overlattice W_f of S_f corresponds to an overlattice M of $L(\Sigma)$, which satisfies the conditions (D2-a)-(D2-c) by Lemma 2.5 and Corollary 2.6. Conversely, suppose that (Σ, MW, T) satisfies the conditions (D1) and (D2). By Lemma 2.4, the condition (D2-b) and $D_H = 0$ imply that there exists an even unimodular overlattice of $M \oplus H \oplus T$ into which $M \oplus H$ and T are primitively embedded. By the theorem of Milnor (see, for example, Serre[15]) on the classification of even unimodular lattices, any even unimodular lattice of

signature $(3, 19)$ is isomorphic to the $K3$ lattice Λ . Then Lemma 2.7 implies that there exists an elliptic $K3$ surface $f : X \rightarrow \mathbb{P}^1$ satisfying $W_f \cong M$ and $T_X \cong T$. The condition $(D2 - c)$ implies $M_{root} = L(\Sigma)$. Combining this with Lemma 2.5, we see that $S_f \cong L(\Sigma)$. Then (2.2) implies that $\Sigma_f = \Sigma$. Using Lemma 2.5 and the condition $(D2 - a)$, we see that $MW_f \cong MW$. Thus the data of $f : X \rightarrow \mathbb{P}^1$ coincides with (Σ, MW, T) . \square

Remark 2.9. In the light of Lemma 2.3, the condition $(D2)$ is equivalent to the following:

$(D3)$ There exists an isotropic subgroup A of $(D_{L(\Sigma)}, q_{L(\Sigma)})$ satisfying the following:

$(D3 - a)$ A is isomorphic to MW ,

$(D3 - b)$ there exists an isomorphism $\gamma : A^\perp/A \xrightarrow{\sim} D_T$ of abelian groups such that $\gamma^*q_T = -q_{L(\Sigma)}|_{A^\perp/A}$, and

$(D3 - c)$ $\text{root}(\phi_{L(\Sigma)}^{-1}(A))$ is equal to $\text{root}(L(\Sigma))$, where $\phi_{L(\Sigma)} : L(\Sigma)^\vee \rightarrow D_{L(\Sigma)}$ is the natural projection.

Remark 2.10. We did not use the conditions $\text{length}(MW) \leq 2$ and $eu(\Sigma) \leq 24$ in the proof of the “if” part of Proposition 2.8. It follows that, if (Σ, MW, T) satisfies $\text{rank}(L(\Sigma)) = 18$ and the condition $(D2)$, then $\text{length}(MW) \leq 2$ and $eu(\Sigma) \leq 24$ follow automatically. This fact can be used when we check the computer program described in the next section.

3. MAKING THE LIST

First we list up all root types Σ satisfying $\text{rank}(L(\Sigma)) = 18$ and $eu(\Sigma) \leq 24$. This list \mathcal{L} consists of 712 elements.

Next we run a program that takes an element Σ of the list \mathcal{L} as an input and proceeds as follows.

Step 1. The program calculates the intersection matrix of $L(\Sigma)^\vee$. Using this matrix, it calculates the discriminant form of $L(\Sigma)$, and decomposes it into p -parts;

$$(D_{L(\Sigma)}, q_{L(\Sigma)}) = \bigoplus_p (D_{L(\Sigma)}, q_{L(\Sigma)})_p,$$

where p runs through the set $\{p_1, \dots, p_k\}$ of prime divisors of the discriminant $|D_{L(\Sigma)}|$ of $L(\Sigma)$. We write the p_i -part of $(D_{L(\Sigma)}, q_{L(\Sigma)})$ by $(D_{L(\Sigma), i}, q_{L(\Sigma), i})$.

Step 2. For each p_i , it calculates the set $I(p_i)$ of all pairs (A, A^\perp) of an isotropic subgroup A of $(D_{L(\Sigma), i}, q_{L(\Sigma), i})$ and its orthogonal complement A^\perp such that $\text{length}(A) \leq 2$.

Step 3. For each element

$$\mathcal{A} := ((A_1, A_1^\perp), \dots, (A_k, A_k^\perp)) \in I(p_1) \times \dots \times I(p_k),$$

it calculates the $\mathbb{Q}/2\mathbb{Z}$ -valued quadratic form

$$q_{\mathcal{A}} := q_{L(\Sigma), 1}|_{A_1^\perp/A_1} \times \dots \times q_{L(\Sigma), k}|_{A_k^\perp/A_k}$$

on the finite abelian group

$$D_{\mathcal{A}} := A_1^\perp/A_1 \times \dots \times A_k^\perp/A_k.$$

Let $d(\mathcal{A})$ be the order of $D_{\mathcal{A}}$.

Step 4. It generates the list $\mathcal{T}(d(\mathcal{A}))$ of positive definite even lattices of rank 2 with discriminant equal to $d(\mathcal{A})$. For each $T \in \mathcal{T}(d(\mathcal{A}))$, it calculates the discriminant form of T and decomposes it into p -parts. If D_T is isomorphic to $D_{\mathcal{A}}$ and q_T is isomorphic to $-q_{\mathcal{A}}$, then it proceeds to the next step. Note that the automorphism group of a finite abelian p -group of length ≤ 2 is easily calculated, and hence it is an easy task to check whether two given quadratic forms on the finite abelian p -group of length ≤ 2 are isomorphic or not.

Step 5. It calculates the Gram matrix of the sublattice $\tilde{L}(\mathcal{A})$ of $L(\Sigma)^\vee$ generated by $L(\Sigma) \subset L(\Sigma)^\vee$ and the pull-backs of generators of the subgroups $A_i \subset D_{L(\Sigma),i}$ by the projection $L(\Sigma)^\vee \rightarrow D_{L(\Sigma)} \rightarrow D_{L(\Sigma),i}$. Then it calculates $\text{root}(\tilde{L}(\mathcal{A}))$ by the method described in the subsection 2.2. If $\text{root}(\tilde{L}(\mathcal{A}))$ is equal to $\text{root}(L(\Sigma))$ calculated by (2.2), then it puts out the pair of the finite abelian group

$$MW := A_1 \times \cdots \times A_k$$

and the lattice T .

Then (Σ, MW, T) satisfies the conditions (D1) and (D3), and all triples (Σ, MW, T) satisfying (D1) and (D3) are obtained by this program.

4. FUNDAMENTAL GROUPS OF OPEN $K3$ SURFACES

A simple normal crossing divisor Δ on a $K3$ surface X is said to be an *ADE-configuration of smooth rational curves* if each irreducible component of Δ is a smooth rational curve and the intersection matrix of the irreducible components of Δ is a direct sum of the Cartan matrices of type A_l , D_m or E_n multiplied by -1 . It is known that Δ is an *ADE-configuration of smooth rational curves* if and only if each connected component of Δ can be contracted to a rational double point. We consider the following quite plausible hypothesis. Let Δ be an *ADE-configuration of smooth rational curves* on a $K3$ surface X .

Hypothesis. If $\pi_1^{\text{alg}}(X \setminus \Delta)$ is trivial, then so is $\pi_1(X \setminus \Delta)$.

Here $\pi_1^{\text{alg}}(X \setminus \Delta)$ is the algebraic fundamental group of $X \setminus \Delta$, which is the pro-finite completion of the topological fundamental group $\pi_1(X \setminus \Delta)$.

Proposition 4.1. *Suppose that Hypothesis is true for any ADE-configuration of smooth rational curves on an arbitrary $K3$ surface. Let Δ be an ADE-configuration of smooth rational curves on a $K3$ surface X . Then $\pi_1(X \setminus \Delta)$ satisfies one of the following:*

- (i) $\pi_1(X \setminus \Delta)$ is trivial.
- (ii) *There exist a complex torus T of dimension 2 and a finite automorphism group G of T such that T/G is birational to X and that $\pi_1(X \setminus \Delta)$ fits in the exact sequence*

$$1 \longrightarrow \pi_1(T) \longrightarrow \pi_1(X \setminus \Delta) \longrightarrow G \longrightarrow 1.$$

- (iii) $\pi_1(X \setminus \Delta)$ is isomorphic to a symplectic automorphism group of a $K3$ surface.

Remark 4.2. Fujiki[4] classified the automorphism groups of complex tori of dimension 2. In particular, the G in (ii) is either one of $\mathbb{Z}/(n)$ ($n = 2, 3, 4, 6$), Q_8 (Quaternion of order 8), D_{12} (Dihedral of order 12) and T_{24} (Tetrahedral of order 24), whence the $\pi_1(X \setminus \Delta)$ in (ii) is a soluble group. Mukai[9] presented the complete list of symplectic automorphism groups of $K3$ surfaces. (See also Kondō[6])

and Xiao[18].) Under Hypothesis, therefore, we know what groups can appear as $\pi_1(X \setminus \Delta)$.

Proof of Proposition 4.1. Suppose that $\pi_1(X \setminus \Delta)$ is non-trivial. By Hypothesis, $\pi_1^{alg}(X \setminus \Delta)$ is also non-trivial. For a surjective homomorphism $\phi : \pi_1(X \setminus \Delta) \rightarrow G$ from $\pi_1(X \setminus \Delta)$ to a finite group G , we denote by

$$\psi_\phi : \tilde{Y}_\phi \longrightarrow X$$

the finite Galois cover of X corresponding to ϕ , which is étale over $X \setminus \Delta$ and whose Galois group is canonically isomorphic to G . Let $\rho : \tilde{Y}'_\phi \rightarrow \tilde{Y}_\phi$ be the resolution of singularities, and $\gamma : \tilde{Y}'_\phi \rightarrow Y_\phi$ the contraction of (-1) -curves. We denote by Δ_ϕ the union of one-dimensional irreducible components of $\gamma(\rho^{-1}(\psi_\phi^{-1}(\Delta)))$. Then it is easy to see that Y_ϕ is either a $K3$ surface or a complex torus of dimension 2, and that the Galois group G of ψ_ϕ acts on Y_ϕ symplectically. Moreover, Δ_ϕ is an empty set or an ADE -configuration of smooth rational curves. We have an exact sequence

$$1 \longrightarrow \pi_1(Y_\phi \setminus \Delta_\phi) \longrightarrow \pi_1(X \setminus \Delta) \longrightarrow G \longrightarrow 1,$$

because $\pi_1(\tilde{Y}'_\phi \setminus \psi_\phi^{-1}(\Delta))$ is isomorphic to $\pi_1(Y_\phi \setminus \Delta_\phi)$. Suppose that there exists a homomorphism $\phi : \pi_1(X \setminus \Delta) \rightarrow G$ such that Y_ϕ is a complex torus of dimension 2. Then Δ_ϕ is empty, and hence (ii) occurs. Suppose that no complex tori of dimension 2 appear as a finite Galois cover of X branched in Δ . Then any finite quotient group of $\pi_1(X \setminus \Delta)$ must appear in Mukai's list of symplectic automorphism groups of $K3$ surfaces. Because this list consists of finite number of isomorphism classes of finite groups, there exists a maximal finite quotient $\phi_{max} : \pi_1(X \setminus \Delta) \rightarrow G_{max}$ of $\pi_1(X \setminus \Delta)$. Then $\pi_1(Y_{\phi_{max}} \setminus \Delta_{\phi_{max}})$ has no non-trivial finite quotient group, and hence it is trivial by Hypothesis. Thus (iii) occurs. \square

For an ADE -configuration Δ of smooth rational curves on a $K3$ surface X , we denote by $\mathbb{Z}[\Delta]$ the sublattice of $H^2(X; \mathbb{Z})$ generated by the cohomology classes of the irreducible components of Δ , which is isomorphic to a negative definite root lattice of type ADE . We denote by Σ_Δ the root type such that $\mathbb{Z}[\Delta]$ is isomorphic to $L(\Sigma_\Delta)$. Using the list of extremal elliptic $K3$ surfaces, we prove the following theorem. We consider the following conditions on a root type Σ .

- (N1) $\text{rank}(L(\Sigma)) \leq 18$, and
- (N2) $\text{length}(D_{L(\Sigma)}) \leq \min\{\text{rank}(L(\Sigma)), 20 - \text{rank}(L(\Sigma))\}$.

Theorem 4.3. *Suppose that a root type Σ_Δ satisfies the conditions (N1) and (N2). If $\mathbb{Z}[\Delta]$ is primitive in $H^2(X; \mathbb{Z})$ then $\pi_1(X \setminus \Delta)$ is trivial.*

By virtue of Lemma 4.6 below, we can easily derive the following:

Corollary 4.4. *Suppose that Σ satisfies the conditions (N1) and (N2). Then Hypothesis is true for any (X, Δ) with $\Sigma_\Delta = \Sigma$. \square*

Remark 4.5. The conditions (N1) and (N2) come from Nikulin[11, Theorem 1.14.1] (see also Morrison[8, Theorem 2.8]), which gives a sufficient condition for the uniqueness of the primitive embedding of $L(\Sigma)$ into the $K3$ lattice Λ .

First we prepare some lemmas. Let $\overline{\mathbb{Z}[\Delta]}$ be the primitive closure of $\mathbb{Z}[\Delta]$ in $H^2(X; \mathbb{Z})$.

Lemma 4.6 (Xiao[18] Lemma 2). *The dual of the abelianisation of $\pi_1(X \setminus \Delta)$ is canonically isomorphic to $\overline{\mathbb{Z}[\Delta]}/\mathbb{Z}[\Delta]$. In particular, if $\pi_1^{\text{alg}}(X \setminus \Delta)$ is trivial, then $\mathbb{Z}[\Delta]$ is primitive in $H^2(X; \mathbb{Z})$. \square*

Let Γ_1 and Γ_2 be graphs with the set of vertices denoted by $\text{Vert}(\Gamma_1)$ and $\text{Vert}(\Gamma_2)$, respectively. An embedding of Γ_1 into Γ_2 is, by definition, an injection $f : \text{Vert}(\Gamma_1) \rightarrow \text{Vert}(\Gamma_2)$ such that, for any $u, v \in \text{Vert}(\Gamma_1)$, $f(u)$ and $f(v)$ are connected by an edge of Γ_2 if and only if u and v are connected by an edge of Γ_1 .

Let $\Gamma(\Sigma)$ denote the Dynkin graph of Σ .

Lemma 4.7. *Suppose that Σ satisfies the conditions (N1) and (N2). Then there exists Σ' satisfying $\text{rank}(L(\Sigma')) = 18$ and the condition (N2) such that $\Gamma(\Sigma)$ can be embedded in $\Gamma(\Sigma')$.*

Proof. This is checked by listing up all Σ satisfying the conditions (N1) and (N2) using computer. \square

Lemma 4.8. *Let $f : X \rightarrow \mathbb{P}^1$ be an elliptic surface with the zero section O . Suppose that a fiber $f^{-1}(v)$ over $v \in \mathbb{P}^1$ is a singular fiber of type III or IV. Let Ξ be a union of some irreducible components of $f^{-1}(v)$ that does not coincide with the whole fiber $f^{-1}(v)$. If U is a small open disk on \mathbb{P}^1 with the center v , then $f^{-1}(U) \setminus (\Xi \cup (f^{-1}(U) \cap O))$ has an abelian fundamental group.*

Proof. This can be proved easily by the van-Kampen theorem. \square

Lemma 4.9. *Let Σ be satisfying the conditions (N1) and (N2). Suppose that (X, Δ) and (X', Δ') satisfy the following:*

- (a) $\Sigma_\Delta = \Sigma_{\Delta'} = \Sigma$,
- (b) $\overline{\mathbb{Z}[\Delta]} = \mathbb{Z}[\Delta]$ and $\overline{\mathbb{Z}[\Delta']} = \mathbb{Z}[\Delta']$.

Then there exists a connected continuous family (X_t, Δ_t) parameterized by $t \in [0, 1]$ such that $(X_0, \Delta_0) = (X, \Delta)$, $(X_1, \Delta_1) = (X', \Delta')$ and that (X_t, Δ_t) are diffeomorphic to one another. In particular, $\pi_1(X \setminus \Delta)$ is isomorphic to $\pi_1(X' \setminus \Delta')$.

Proof. By Nikulin[11, Theorem 1.14.1], the primitive embedding of $L(\Sigma)$ into the K3 lattice Λ is unique up to $\text{Aut}(\Lambda)$. Hence the assertion follows from Nikulin's connectedness theorem[10, Theorem 2.10]. \square

Proof of Theorem 4.3. Let us consider the following:

Claim 1. *Suppose that Σ satisfies $\text{rank}(L(\Sigma)) = 18$ and the condition (N2). Then there exists an ADE-configuration of smooth rational curves Δ_Σ on a K3 surface X_Σ such that $\Sigma_{\Delta_\Sigma} = \Sigma$ and $\pi_1(X_\Sigma \setminus \Delta_\Sigma) = \{1\}$.*

We deduce Theorem 4.3 from Claim 1. Suppose that Δ is an ADE-configuration of smooth rational curves on a K3 surface X such that Σ_Δ satisfies the conditions (N1) and (N2), and that $\mathbb{Z}[\Delta]$ is primitive in $H^2(X; \mathbb{Z})$. By Lemma 4.7, there exists Σ_1 satisfying $\text{rank}(L(\Sigma_1)) = 18$ and the condition (N2) such that $\Gamma(\Sigma_\Delta)$ is embedded into $\Gamma(\Sigma_1)$. By Claim 1, we have (X_1, Δ_1) such that $\Sigma_{\Delta_1} = \Sigma_1$ and $\pi_1(X_1 \setminus \Delta_1) = \{1\}$. Let $\Delta' \subset \Delta_1$ be the sub-configuration of smooth rational curves on X_1 which corresponds to the subgraph $\Gamma(\Sigma_\Delta) \hookrightarrow \Gamma(\Sigma_1) = \Gamma(\Sigma_{\Delta_1})$. There is a surjection from $\pi_1(X_1 \setminus \Delta_1)$ to $\pi_1(X_1 \setminus \Delta')$, and hence $\pi_1(X_1 \setminus \Delta')$ is trivial. In particular, $\mathbb{Z}[\Delta']$ is primitive in $H^2(X_1; \mathbb{Z})$. Since $\Sigma_{\Delta'} = \Sigma_\Delta$, Lemma 4.9 implies that $\pi_1(X \setminus \Delta)$ is isomorphic to $\pi_1(X_1 \setminus \Delta')$. Thus $\pi_1(X \setminus \Delta)$ is trivial.

Let $f : X \rightarrow \mathbb{P}^1$ be an extremal elliptic $K3$ surface. For a point $v \in R_f$, we denote the total fiber of f over v by

$$\sum_{i=1}^{r_v} m_{v,i} C_{v,i},$$

where $m_{v,i}$ is the multiplicity of the irreducible component $C_{v,i}$ of $f^{-1}(v)$. We denote by Γ_f the union of the zero section and all irreducible fibers $f^{-1}(v)$ ($v \in R_f$).

Claim 2. Suppose that $MW_f = (0)$. Suppose that a sub-configuration Δ of Γ_f satisfies the following two conditions.

(Z1) The number of $v \in R_f$ such that $m_{v,i} = 1 \implies$ The number of $C_{v,i} \subset \Delta$ is at most one.

(Z2) Either one of the following holds:

(Z2-a) The configuration Δ does not contain the zero section,

(Z2-b) there is a point $v_1 \in R_f$ such that the type $\tau(S_{f,v_1})$ is A_1 and that $F_1 := f^{-1}(v_1)$ and Δ have no common irreducible components, or

(Z2-c) $eu(\Sigma_f) \leq 23$.

Then $\pi_1(X \setminus \Delta)$ is trivial.

Proof of Claim 2. By Lemma 2.5, the assumption $MW_f = (0)$ implies that the cohomology classes $[O]$ and $[C_{v,i}]$ ($v \in R_f, i = 1, \dots, r_v$) of the irreducible components of Γ_f span NS_X . The relations among these generators are generated by

$$\sum_{i=1}^{r_v} m_{v,i} C_{v,i} = \sum_{i=1}^{r_{v'}} m_{v',i} C_{v',i} \quad (v, v' \in R_f).$$

Therefore the condition (Z1) implies that the cohomology classes of the irreducible components of Δ constitute a subset of a \mathbb{Z} -basis of NS_X . Hence $\mathbb{Z}[\Delta]$ is primitive in $H^2(X; \mathbb{Z})$. In particular, $\pi_1(X \setminus \Delta)$ is a perfect group by Lemma 4.6. On the other hand, the condition (Z1) implies that there exists a point $v_0 \in \mathbb{P}^1$ such that every fiber of the restriction

$$f|_{X \setminus (\Delta \cup f^{-1}(v_0))} : X \setminus (\Delta \cup f^{-1}(v_0)) \longrightarrow \mathbb{P}^1 \setminus \{v_0\}$$

of f has a reduced irreducible component. Then, by Nori's lemma [13, Lemma 1.5 (C)], if U is a non-empty connected classically open subset of $\mathbb{P}^1 \setminus \{v_0\}$, then the inclusion of $f^{-1}(U) \setminus (f^{-1}(U) \cap \Delta)$ into $X \setminus (\Delta \cup f^{-1}(v_0))$ induces a surjection on the fundamental groups. The inclusion of $X \setminus (\Delta \cup f^{-1}(v_0))$ into $X \setminus \Delta$ also induces a surjection on the fundamental groups. We shall show that there exists a small open disk U on $\mathbb{P}^1 \setminus \{v_0\}$ such that

$$G_U := \pi_1(f^{-1}(U) \setminus (f^{-1}(U) \cap \Delta))$$

is abelian. When (Z2-a) occurs, we take a small open disk disjoint from R_f as U . Then G_U is abelian, because of $f^{-1}(U) \cap \Delta = \emptyset$. Suppose that (Z2-b) occurs. We can take v_0 from $\mathbb{P}^1 \setminus \{v_1\}$, because F_1 has no irreducible components of multiplicity ≥ 2 . We choose a small open disk U with the center v_1 . There is a contraction from $f^{-1}(U) \setminus (f^{-1}(U) \cap \Delta)$ to $F_1 \setminus (F_1 \cap \Delta)$. Because $\pi_1(F_1 \setminus (F_1 \cap \Delta))$ is abelian, so is G_U . Suppose that (Z2-c) occurs. By Lemma 2.2, there exists a singular fiber $F_2 := f^{-1}(v_2)$ of type I_1, II, III or IV . Because F_2 has no irreducible components of multiplicity ≥ 2 , we can choose v_0 from $\mathbb{P}^1 \setminus \{v_2\}$. If F_2 is of type I_1 or II , then $F_2 \cap \Delta$ consists of a nonsingular point of F_2 , and $\pi_1(F_2 \setminus (F_2 \cap \Delta))$ is abelian. Hence

G_U is also abelian. If F_2 is of type III or IV, then $F_2 \cap \Delta$ cannot coincide with the whole fiber F_2 . Hence Lemma 4.8 implies that G_U is abelian. Therefore we see that $\pi_1(X \setminus \Delta)$ is abelian. Being both perfect and abelian, $\pi_1(X \setminus \Delta)$ is trivial. \square

Now we proceed to the proof of Claim 1. We list up all Σ satisfying the condition (N2) and $\text{rank}(L(\Sigma)) = 18$. It consists of 297 elements. Among them, 199 elements can be the type Σ_f of singular fibers of some extremal elliptic $K3$ surface $f : X \rightarrow \mathbb{P}^1$ with $MW_f = 0$. For these configurations, $\pi_1(X \setminus \Delta)$ is trivial by Claim 2. The remaining 98 configurations are listed in the second column of Table 1 below. Each of them is a sub-configuration of Γ_f satisfying the conditions (Z1) and (Z2), where $f : X \rightarrow \mathbb{P}^1$ is the extremal elliptic $K3$ surface with $MW_f = 0$ whose number in Table 2 is given in the third column of Table 1. The fourth and fifth columns of Table 1 indicate Σ_f and $eu(\Sigma_f)$, respectively. In the case nos. 20, 28, 39, 41 and 85 in Table 1, we can choose the embedding of Δ into Γ_f in such a way that (Z2 - b) holds. In the case nos. 30, 37, 57 and 63 in Table 1, we can choose the embedding of Δ into Γ_f in such a way that (Z2 - a) holds. By Claim 2 again, $\pi_1(X \setminus \Delta)$ is trivial for these 98 configurations Δ . \square

Remark 4.10. The graph $\Gamma(A_{19})$ (resp. $\Gamma(D_{19})$) can be embedded into Γ_f in such a way that (Z1) and (Z2) are satisfied, where $f : X \rightarrow \mathbb{P}^1$ is the extremal elliptic $K3$ surfaces whose number in Table 2 is 312 (resp. 320). Therefore, if $\Gamma(\Delta)$ is embedded in $\Gamma(A_{19})$ or $\Gamma(D_{19})$, then $\Gamma(\Delta)$ can be embedded in Γ_f in such a way that (Z1) and (Z2) are satisfied.

Table 1. List of embedding of Δ in Γ_f

| no | Δ | No | Σ_f | $eu(\Sigma_f)$ |
|----|-----------------------------|-----|---------------------------------|----------------|
| 1 | $A_2 + A_3 + 2 A_4 + A_5$ | 19 | $A_2 + 2 A_3 + A_4 + A_6$ | 23 |
| 2 | $A_1 + A_2 + A_3 + 2 A_6$ | 23 | $A_1 + A_2 + A_4 + A_5 + A_6$ | 23 |
| 3 | $2 A_1 + A_4 + 2 A_6$ | 23 | $A_1 + A_2 + A_4 + A_5 + A_6$ | 23 |
| 4 | $2 A_2 + 2 A_4 + A_6$ | 23 | $A_1 + A_2 + A_4 + A_5 + A_6$ | 23 |
| 5 | $A_1 + A_5 + 2 A_6$ | 40 | $A_1 + A_4 + A_6 + A_7$ | 22 |
| 6 | $A_4 + 2 A_7$ | 52 | $A_4 + A_6 + A_8$ | 21 |
| 7 | $A_1 + A_2 + 2 A_4 + A_7$ | 23 | $A_1 + A_2 + A_4 + A_5 + A_6$ | 23 |
| 8 | $A_3 + 2 A_4 + A_7$ | 24 | $A_3 + A_4 + A_5 + A_6$ | 22 |
| 9 | $A_2 + 2 A_4 + A_8$ | 36 | $A_2 + A_4 + A_5 + A_7$ | 22 |
| 10 | $2 A_3 + A_4 + A_8$ | 46 | $A_1 + A_2 + A_3 + A_4 + A_8$ | 23 |
| 11 | $A_3 + A_7 + A_8$ | 53 | $A_1 + A_2 + A_7 + A_8$ | 22 |
| 12 | $A_1 + 2 A_2 + A_4 + A_9$ | 46 | $A_1 + A_2 + A_3 + A_4 + A_8$ | 23 |
| 13 | $A_2 + A_3 + A_4 + A_9$ | 71 | $2 A_2 + A_4 + A_{10}$ | 22 |
| 14 | $A_3 + A_4 + A_{11}$ | 93 | $A_2 + A_4 + A_{12}$ | 21 |
| 15 | $A_7 + A_{11}$ | 312 | $A_{10} + E_8$ | 21 |
| 16 | $2 A_3 + A_{12}$ | 93 | $A_2 + A_4 + A_{12}$ | 21 |
| 17 | $A_3 + A_{15}$ | 312 | $A_{10} + E_8$ | 21 |
| 18 | $A_2 + 2 A_6 + D_4$ | 99 | $A_2 + A_3 + A_{13}$ | 21 |
| 19 | $2 A_4 + A_6 + D_4$ | 18 | $A_1 + A_3 + 2 A_4 + A_6$ | 23 |
| 20 | $2 A_2 + A_4 + A_6 + D_4$ | 20 | $A_1 + 2 A_2 + A_3 + A_4 + A_6$ | 24 |
| 21 | $A_2 + A_4 + A_8 + D_4$ | 44 | $2 A_1 + 2 A_4 + A_8$ | 23 |
| 22 | $A_6 + A_8 + D_4$ | 50 | $2 A_1 + A_2 + A_6 + A_8$ | 23 |
| 23 | $2 A_2 + A_{10} + D_4$ | 72 | $2 A_1 + A_2 + A_4 + A_{10}$ | 23 |
| 24 | $A_4 + A_{10} + D_4$ | 72 | $2 A_1 + A_2 + A_4 + A_{10}$ | 23 |
| 25 | $A_2 + A_{12} + D_4$ | 90 | $2 A_1 + 2 A_2 + A_{12}$ | 23 |
| 26 | $A_{14} + D_4$ | 320 | $D_{10} + E_8$ | 22 |
| 27 | $2 A_2 + A_4 + 2 D_5$ | 210 | $2 A_2 + D_{14}$ | 22 |
| 28 | $A_1 + 2 A_2 + 2 A_4 + D_5$ | 157 | $A_1 + A_2 + 2 A_4 + D_7$ | 24 |
| 29 | $A_2 + A_3 + 2 A_4 + D_5$ | 46 | $A_1 + A_2 + A_3 + A_4 + A_8$ | 23 |
| 30 | $A_2 + A_6 + 2 D_5$ | 193 | $A_2 + A_6 + D_{10}$ | 22 |
| 31 | $A_3 + A_4 + A_6 + D_5$ | 18 | $A_1 + A_3 + 2 A_4 + A_6$ | 23 |
| 32 | $A_2 + A_4 + A_7 + D_5$ | 72 | $2 A_1 + A_2 + A_4 + A_{10}$ | 23 |
| 33 | $A_6 + A_7 + D_5$ | 50 | $2 A_1 + A_2 + A_6 + A_8$ | 23 |
| 34 | $A_2 + A_3 + A_8 + D_5$ | 50 | $2 A_1 + A_2 + A_6 + A_8$ | 23 |
| 35 | $A_3 + A_{10} + D_5$ | 69 | $A_1 + 2 A_2 + A_3 + A_{10}$ | 23 |
| 36 | $A_2 + A_{11} + D_5$ | 90 | $2 A_1 + 2 A_2 + A_{12}$ | 23 |
| 37 | $A_4 + 2 D_7$ | 213 | $A_4 + D_{14}$ | 21 |
| 38 | $A_3 + 2 A_4 + D_7$ | 44 | $2 A_1 + 2 A_4 + A_8$ | 23 |
| 39 | $2 A_2 + A_3 + A_4 + D_7$ | 20 | $A_1 + 2 A_2 + A_3 + A_4 + A_6$ | 24 |
| 40 | $A_2 + A_4 + A_5 + D_7$ | 23 | $A_1 + A_2 + A_4 + A_5 + A_6$ | 23 |
| 41 | $A_1 + 2 A_2 + A_6 + D_7$ | 14 | $2 A_1 + 2 A_2 + 2 A_6$ | 24 |

Table 1. List of embedding of Δ in Γ_f

| no | Δ | No | Σ_f | $eu(\Sigma_f)$ |
|----|-------------------------------|-----|-------------------------------|----------------|
| 42 | $2 A_2 + A_7 + D_7$ | 90 | $2 A_1 + 2 A_2 + A_{12}$ | 23 |
| 43 | $A_4 + A_7 + D_7$ | 44 | $2 A_1 + 2 A_4 + A_8$ | 23 |
| 44 | $A_1 + A_2 + A_8 + D_7$ | 50 | $2 A_1 + A_2 + A_6 + A_8$ | 23 |
| 45 | $A_3 + A_8 + D_7$ | 44 | $2 A_1 + 2 A_4 + A_8$ | 23 |
| 46 | $A_{11} + D_7$ | 320 | $D_{10} + E_8$ | 22 |
| 47 | $A_2 + A_4 + D_5 + D_7$ | 200 | $A_2 + A_5 + D_{11}$ | 22 |
| 48 | $A_6 + D_5 + D_7$ | 186 | $A_9 + D_9$ | 21 |
| 49 | $A_2 + 2 A_4 + D_8$ | 66 | $A_2 + A_7 + A_9$ | 21 |
| 50 | $A_4 + A_6 + D_8$ | 23 | $A_1 + A_2 + A_4 + A_5 + A_6$ | 23 |
| 51 | $A_2 + A_8 + D_8$ | 50 | $2 A_1 + A_2 + A_6 + A_8$ | 23 |
| 52 | $A_{10} + D_8$ | 320 | $D_{10} + E_8$ | 22 |
| 53 | $A_1 + 2 A_4 + D_9$ | 44 | $2 A_1 + 2 A_4 + A_8$ | 23 |
| 54 | $A_2 + A_3 + A_4 + D_9$ | 46 | $A_1 + A_2 + A_3 + A_4 + A_8$ | 23 |
| 55 | $A_3 + A_6 + D_9$ | 76 | $2 A_1 + A_6 + A_{10}$ | 22 |
| 56 | $A_2 + A_7 + D_9$ | 50 | $2 A_1 + A_2 + A_6 + A_8$ | 23 |
| 57 | $2 A_2 + D_5 + D_9$ | 210 | $2 A_2 + D_{14}$ | 22 |
| 58 | $A_2 + D_7 + D_9$ | 186 | $A_9 + D_9$ | 21 |
| 59 | $2 A_2 + A_4 + D_{10}$ | 72 | $2 A_1 + A_2 + A_4 + A_{10}$ | 23 |
| 60 | $A_3 + A_4 + D_{11}$ | 44 | $2 A_1 + 2 A_4 + A_8$ | 23 |
| 61 | $A_7 + D_{11}$ | 320 | $D_{10} + E_8$ | 22 |
| 62 | $A_2 + D_5 + D_{11}$ | 186 | $A_9 + D_9$ | 21 |
| 63 | $D_7 + D_{11}$ | 218 | D_{18} | 20 |
| 64 | $A_2 + A_4 + D_{12}$ | 72 | $2 A_1 + A_2 + A_4 + A_{10}$ | 23 |
| 65 | $A_6 + D_{12}$ | 320 | $D_{10} + E_8$ | 22 |
| 66 | $A_1 + 2 A_2 + D_{13}$ | 90 | $2 A_1 + 2 A_2 + A_{12}$ | 23 |
| 67 | $A_2 + A_3 + D_{13}$ | 72 | $2 A_1 + A_2 + A_4 + A_{10}$ | 23 |
| 68 | $A_3 + D_{15}$ | 320 | $D_{10} + E_8$ | 22 |
| 69 | $A_2 + D_{16}$ | 320 | $D_{10} + E_8$ | 22 |
| 70 | $2 A_1 + A_4 + 2 E_6$ | 303 | $A_1 + A_4 + A_5 + E_8$ | 23 |
| 71 | $2 A_1 + A_2 + 2 A_4 + E_6$ | 23 | $A_1 + A_2 + A_4 + A_5 + A_6$ | 23 |
| 72 | $A_2 + 2 A_3 + A_4 + E_6$ | 46 | $A_1 + A_2 + A_3 + A_4 + A_8$ | 23 |
| 73 | $2 A_6 + E_6$ | 37 | $A_1 + 2 A_2 + A_6 + A_7$ | 23 |
| 74 | $2 A_3 + A_6 + E_6$ | 41 | $A_5 + A_6 + A_7$ | 21 |
| 75 | $A_2 + A_3 + A_7 + E_6$ | 37 | $A_1 + 2 A_2 + A_6 + A_7$ | 23 |
| 76 | $2 A_4 + D_4 + E_6$ | 182 | $A_4 + A_5 + D_9$ | 22 |
| 77 | $A_2 + A_6 + D_4 + E_6$ | 183 | $A_1 + A_2 + A_6 + D_9$ | 23 |
| 78 | $A_8 + D_4 + E_6$ | 186 | $A_9 + D_9$ | 21 |
| 79 | $A_1 + D_5 + 2 E_6$ | 320 | $D_{10} + E_8$ | 22 |
| 80 | $A_2 + 2 D_5 + E_6$ | 320 | $D_{10} + E_8$ | 22 |
| 81 | $A_1 + A_2 + A_4 + D_5 + E_6$ | 193 | $A_2 + A_6 + D_{10}$ | 22 |
| 82 | $A_2 + A_3 + D_7 + E_6$ | 200 | $A_2 + A_5 + D_{11}$ | 22 |

Table 1. List of embedding of Δ in Γ_f

| no | Δ | No | Σ_f | $eu(\Sigma_f)$ |
|----|-----------------------------|-----|-------------------------------|----------------|
| 83 | $A_5 + D_7 + E_6$ | 320 | $D_{10} + E_8$ | 22 |
| 84 | $A_2 + D_{10} + E_6$ | 193 | $A_2 + A_6 + D_{10}$ | 22 |
| 85 | $A_1 + A_2 + 2 A_4 + E_7$ | 17 | $2 A_1 + A_2 + 2 A_4 + A_6$ | 24 |
| 86 | $A_3 + 2 A_4 + E_7$ | 18 | $A_1 + A_3 + 2 A_4 + A_6$ | 23 |
| 87 | $2 A_2 + D_7 + E_7$ | 210 | $2 A_2 + D_{14}$ | 22 |
| 88 | $A_2 + 2 A_4 + E_8$ | 36 | $A_2 + A_4 + A_5 + A_7$ | 22 |
| 89 | $2 A_1 + 2 A_2 + A_4 + E_8$ | 30 | $2 A_2 + A_3 + A_4 + A_7$ | 23 |
| 90 | $2 A_3 + A_4 + E_8$ | 24 | $A_3 + A_4 + A_5 + A_6$ | 22 |
| 91 | $A_3 + A_7 + E_8$ | 46 | $A_1 + A_2 + A_3 + A_4 + A_8$ | 23 |
| 92 | $A_2 + A_4 + D_4 + E_8$ | 182 | $A_4 + A_5 + D_9$ | 22 |
| 93 | $A_6 + D_4 + E_8$ | 186 | $A_9 + D_9$ | 21 |
| 94 | $A_1 + 2 A_2 + D_5 + E_8$ | 210 | $2 A_2 + D_{14}$ | 22 |
| 95 | $A_2 + A_3 + D_5 + E_8$ | 198 | $2 A_2 + A_3 + D_{11}$ | 23 |
| 96 | $A_3 + D_7 + E_8$ | 213 | $A_4 + D_{14}$ | 21 |
| 97 | $A_2 + D_8 + E_8$ | 210 | $2 A_2 + D_{14}$ | 22 |
| 98 | $2 A_1 + A_2 + E_6 + E_8$ | 320 | $D_{10} + E_8$ | 22 |

Table 2. List of extremal elliptic $K3$ surfaces

| No | Σ | MW | a | b | c |
|----|--------------------------------|--|-----|-----|-----|
| 1 | $6A_3$ | $\mathbb{Z}/(4) \times \mathbb{Z}/(4)$ | 4 | 0 | 4 |
| 2 | $2A_1 + 4A_4$ | $\mathbb{Z}/(5)$ | 10 | 0 | 10 |
| 3 | $2A_2 + 2A_3 + 2A_4$ | (0) | 60 | 0 | 60 |
| 4 | $3A_1 + 3A_5$ | $\mathbb{Z}/(2) \times \mathbb{Z}/(6)$ | 2 | 0 | 6 |
| 5 | $4A_2 + 2A_5$ | $\mathbb{Z}/(3) \times \mathbb{Z}/(3)$ | 6 | 0 | 6 |
| 6 | $A_3 + 3A_5$ | $\mathbb{Z}/(6)$ | 4 | 0 | 6 |
| 7 | $2A_1 + 2A_3 + 2A_5$ | $\mathbb{Z}/(2) \times \mathbb{Z}/(2)$ | 12 | 0 | 12 |
| 8 | $A_1 + 2A_2 + A_3 + 2A_5$ | $\mathbb{Z}/(6)$ | 6 | 0 | 12 |
| 9 | $2A_4 + 2A_5$ | (0) | 30 | 0 | 30 |
| 10 | $2A_2 + A_4 + 2A_5$ | $\mathbb{Z}/(3)$ | 6 | 0 | 30 |
| 11 | $A_1 + A_3 + A_4 + 2A_5$ | $\mathbb{Z}/(2)$ | 12 | 0 | 30 |
| 12 | $A_1 + A_2 + 2A_3 + A_4 + A_5$ | $\mathbb{Z}/(2)$ | 24 | 12 | 36 |
| 13 | $3A_6$ | $\mathbb{Z}/(7)$ | 2 | 1 | 4 |
| 14 | $2A_1 + 2A_2 + 2A_6$ | (0) | 42 | 0 | 42 |
| 15 | $2A_3 + 2A_6$ | (0) | 28 | 0 | 28 |
| 16 | $A_2 + A_4 + 2A_6$ | (0) | 28 | 7 | 28 |
| 17 | $2A_1 + A_2 + 2A_4 + A_6$ | (0) | 50 | 20 | 50 |
| 18 | $A_1 + A_3 + 2A_4 + A_6$ | (0) | 10 | 0 | 140 |
| | | | 20 | 0 | 70 |
| 19 | $A_2 + 2A_3 + A_4 + A_6$ | (0) | 24 | 12 | 76 |
| 20 | $A_1 + 2A_2 + A_3 + A_4 + A_6$ | (0) | 30 | 0 | 84 |
| 21 | $2A_1 + 2A_5 + A_6$ | $\mathbb{Z}/(2)$ | 12 | 6 | 24 |
| 22 | $A_1 + 2A_3 + A_5 + A_6$ | $\mathbb{Z}/(2)$ | 4 | 0 | 84 |
| 23 | $A_1 + A_2 + A_4 + A_5 + A_6$ | (0) | 30 | 0 | 42 |
| | | | 18 | 6 | 72 |
| 24 | $A_3 + A_4 + A_5 + A_6$ | (0) | 12 | 0 | 70 |
| 25 | $4A_1 + 2A_7$ | $\mathbb{Z}/(2) \times \mathbb{Z}/(4)$ | 4 | 0 | 4 |
| 26 | $2A_2 + 2A_7$ | (0) | 24 | 0 | 24 |
| | | $\mathbb{Z}/(2)$ | 12 | 0 | 12 |
| 27 | $A_1 + A_3 + 2A_7$ | $\mathbb{Z}/(8)$ | 2 | 0 | 4 |
| 28 | $2A_1 + 3A_3 + A_7$ | $\mathbb{Z}/(2) \times \mathbb{Z}/(4)$ | 4 | 0 | 8 |
| 29 | $A_2 + 3A_3 + A_7$ | $\mathbb{Z}/(4)$ | 4 | 0 | 24 |
| 30 | $2A_2 + A_3 + A_4 + A_7$ | (0) | 12 | 0 | 120 |
| 31 | $2A_1 + A_2 + A_3 + A_4 + A_7$ | $\mathbb{Z}/(2)$ | 20 | 0 | 24 |
| 32 | $A_1 + 2A_5 + A_7$ | $\mathbb{Z}/(2)$ | 6 | 0 | 24 |
| 33 | $3A_1 + A_3 + A_5 + A_7$ | $\mathbb{Z}/(2) \times \mathbb{Z}/(2)$ | 8 | 0 | 12 |

Table 2. List of extremal elliptic $K3$ surfaces

| No | Σ | MW | a | b | c |
|----|-------------------------------|------------------|-----|-----|-----|
| 34 | $A_1 + A_2 + A_3 + A_5 + A_7$ | $\mathbb{Z}/(2)$ | 12 | 0 | 24 |
| 35 | $2A_1 + A_4 + A_5 + A_7$ | $\mathbb{Z}/(2)$ | 2 | 0 | 120 |
| 36 | $A_2 + A_4 + A_5 + A_7$ | (0) | 6 | 0 | 120 |
| | | | 24 | 0 | 30 |
| 37 | $A_1 + 2A_2 + A_6 + A_7$ | (0) | 24 | 0 | 42 |
| 38 | $2A_1 + A_3 + A_6 + A_7$ | $\mathbb{Z}/(2)$ | 12 | 4 | 20 |
| 39 | $A_2 + A_3 + A_6 + A_7$ | (0) | 4 | 0 | 168 |
| 40 | $A_1 + A_4 + A_6 + A_7$ | (0) | 2 | 0 | 280 |
| | | | 18 | 4 | 32 |
| 41 | $A_5 + A_6 + A_7$ | (0) | 16 | 4 | 22 |
| 42 | $2A_1 + 2A_8$ | (0) | 18 | 0 | 18 |
| | | $\mathbb{Z}/(3)$ | 4 | 2 | 10 |
| 43 | $A_1 + 3A_2 + A_3 + A_8$ | $\mathbb{Z}/(3)$ | 12 | 0 | 18 |
| 44 | $2A_1 + 2A_4 + A_8$ | (0) | 20 | 10 | 50 |
| 45 | $3A_2 + A_4 + A_8$ | $\mathbb{Z}/(3)$ | 12 | 3 | 12 |
| 46 | $A_1 + A_2 + A_3 + A_4 + A_8$ | (0) | 6 | 0 | 180 |
| 47 | $A_1 + 2A_2 + A_5 + A_8$ | $\mathbb{Z}/(3)$ | 6 | 0 | 18 |
| 48 | $A_2 + A_3 + A_5 + A_8$ | $\mathbb{Z}/(3)$ | 4 | 0 | 18 |
| 49 | $A_1 + A_4 + A_5 + A_8$ | (0) | 18 | 0 | 30 |
| 50 | $2A_1 + A_2 + A_6 + A_8$ | (0) | 18 | 0 | 42 |
| 51 | $A_1 + A_3 + A_6 + A_8$ | (0) | 10 | 4 | 52 |
| 52 | $A_4 + A_6 + A_8$ | (0) | 18 | 9 | 22 |
| 53 | $A_1 + A_2 + A_7 + A_8$ | (0) | 18 | 0 | 24 |
| 54 | $2A_9$ | (0) | 10 | 0 | 10 |
| | | $\mathbb{Z}/(5)$ | 2 | 0 | 2 |
| 55 | $A_1 + A_2 + 2A_3 + A_9$ | $\mathbb{Z}/(2)$ | 4 | 0 | 60 |
| 56 | $2A_1 + 2A_2 + A_3 + A_9$ | $\mathbb{Z}/(2)$ | 6 | 0 | 60 |
| 57 | $A_1 + 2A_4 + A_9$ | $\mathbb{Z}/(5)$ | 2 | 0 | 10 |
| 58 | $3A_1 + A_2 + A_4 + A_9$ | $\mathbb{Z}/(2)$ | 20 | 10 | 20 |
| 59 | $2A_1 + A_3 + A_4 + A_9$ | $\mathbb{Z}/(2)$ | 10 | 0 | 20 |
| 60 | $2A_1 + A_2 + A_5 + A_9$ | $\mathbb{Z}/(2)$ | 12 | 6 | 18 |
| 61 | $A_1 + A_3 + A_5 + A_9$ | $\mathbb{Z}/(2)$ | 10 | 0 | 12 |
| 62 | $A_4 + A_5 + A_9$ | (0) | 10 | 0 | 30 |
| | | $\mathbb{Z}/(2)$ | 10 | 5 | 10 |
| 63 | $3A_1 + A_6 + A_9$ | $\mathbb{Z}/(2)$ | 4 | 2 | 36 |
| 64 | $A_1 + A_2 + A_6 + A_9$ | (0) | 10 | 0 | 42 |

Table 2. List of extremal elliptic $K3$ surfaces

| No | Σ | MW | a | b | c |
|----|-----------------------------|------------------|-----|-----|-----|
| 65 | $A_3 + A_6 + A_9$ | (0) | 2 | 0 | 140 |
| 66 | $A_2 + A_7 + A_9$ | (0) | 10 | 0 | 24 |
| 67 | $A_1 + A_8 + A_9$ | (0) | 10 | 0 | 18 |
| 68 | $A_2 + 2A_3 + A_{10}$ | (0) | 24 | 12 | 28 |
| 69 | $A_1 + 2A_2 + A_3 + A_{10}$ | (0) | 12 | 0 | 66 |
| 70 | $2A_4 + A_{10}$ | (0) | 10 | 5 | 30 |
| 71 | $2A_2 + A_4 + A_{10}$ | (0) | 6 | 3 | 84 |
| | | | 24 | 9 | 24 |
| 72 | $2A_1 + A_2 + A_4 + A_{10}$ | (0) | 2 | 0 | 330 |
| 73 | $A_1 + A_3 + A_4 + A_{10}$ | (0) | 20 | 0 | 22 |
| | | | 12 | 4 | 38 |
| 74 | $A_1 + A_2 + A_5 + A_{10}$ | (0) | 6 | 0 | 66 |
| | | | 18 | 6 | 24 |
| 75 | $A_3 + A_5 + A_{10}$ | (0) | 4 | 0 | 66 |
| | | | 12 | 0 | 22 |
| 76 | $2A_1 + A_6 + A_{10}$ | (0) | 12 | 2 | 26 |
| 77 | $A_2 + A_6 + A_{10}$ | (0) | 4 | 1 | 58 |
| | | | 16 | 5 | 16 |
| 78 | $A_1 + A_7 + A_{10}$ | (0) | 2 | 0 | 88 |
| | | | 10 | 2 | 18 |
| 79 | $A_8 + A_{10}$ | (0) | 10 | 1 | 10 |
| 80 | $A_1 + 3A_2 + A_{11}$ | $\mathbb{Z}/(3)$ | 6 | 0 | 12 |
| 81 | $3A_1 + 2A_2 + A_{11}$ | $\mathbb{Z}/(6)$ | 2 | 0 | 12 |
| 82 | $A_1 + 2A_3 + A_{11}$ | $\mathbb{Z}/(4)$ | 4 | 0 | 6 |
| 83 | $2A_2 + A_3 + A_{11}$ | $\mathbb{Z}/(3)$ | 4 | 0 | 12 |
| | | $\mathbb{Z}/(6)$ | 4 | 2 | 4 |
| 84 | $2A_1 + A_2 + A_3 + A_{11}$ | $\mathbb{Z}/(4)$ | 6 | 0 | 6 |
| | | $\mathbb{Z}/(2)$ | 12 | 0 | 12 |
| 85 | $3A_1 + A_4 + A_{11}$ | $\mathbb{Z}/(2)$ | 6 | 0 | 20 |
| 86 | $A_1 + A_2 + A_4 + A_{11}$ | (0) | 12 | 0 | 30 |
| 87 | $2A_1 + A_5 + A_{11}$ | $\mathbb{Z}/(2)$ | 6 | 0 | 12 |
| | | $\mathbb{Z}/(6)$ | 2 | 0 | 4 |
| 88 | $A_2 + A_5 + A_{11}$ | $\mathbb{Z}/(3)$ | 4 | 0 | 6 |
| 89 | $A_1 + A_6 + A_{11}$ | (0) | 4 | 0 | 42 |
| 90 | $2A_1 + 2A_2 + A_{12}$ | (0) | 12 | 6 | 42 |
| 91 | $A_1 + A_2 + A_3 + A_{12}$ | (0) | 6 | 0 | 52 |

Table 2. List of extremal elliptic $K3$ surfaces

| No | Σ | MW | a | b | c |
|-----|-------------------------------|------------------|------------------|-----|-----|
| 92 | $2A_1 + A_4 + A_{12}$ | (0) | 2 | 0 | 130 |
| | | | 18 | 8 | 18 |
| 93 | $A_2 + A_4 + A_{12}$ | (0) | 6 | 3 | 34 |
| 94 | $A_1 + A_5 + A_{12}$ | (0) | 10 | 2 | 16 |
| 95 | $A_6 + A_{12}$ | (0) | 2 | 1 | 46 |
| 96 | $A_1 + 2A_2 + A_{13}$ | (0) | 6 | 0 | 42 |
| | | | $\mathbb{Z}/(2)$ | 6 | 3 |
| 97 | $3A_1 + A_2 + A_{13}$ | $\mathbb{Z}/(2)$ | 2 | 0 | 42 |
| 98 | $2A_1 + A_3 + A_{13}$ | $\mathbb{Z}/(2)$ | 6 | 2 | 10 |
| 99 | $A_2 + A_3 + A_{13}$ | (0) | 4 | 0 | 42 |
| 100 | $A_1 + A_4 + A_{13}$ | (0) | 2 | 0 | 70 |
| | | | 8 | 2 | 18 |
| | | | $\mathbb{Z}/(2)$ | 2 | 1 |
| 101 | $A_5 + A_{13}$ | (0) | 4 | 2 | 22 |
| 102 | $2A_2 + A_{14}$ | $\mathbb{Z}/(3)$ | 4 | 1 | 4 |
| 103 | $2A_1 + A_2 + A_{14}$ | (0) | 12 | 6 | 18 |
| | | | $\mathbb{Z}/(3)$ | 2 | 0 |
| 104 | $A_1 + A_3 + A_{14}$ | (0) | 10 | 0 | 12 |
| 105 | $A_4 + A_{14}$ | (0) | 10 | 5 | 10 |
| 106 | $3A_1 + A_{15}$ | $\mathbb{Z}/(4)$ | 2 | 0 | 4 |
| 107 | $A_1 + A_2 + A_{15}$ | (0) | 10 | 2 | 10 |
| | | | $\mathbb{Z}/(2)$ | 4 | 0 |
| 108 | $A_3 + A_{15}$ | $\mathbb{Z}/(4)$ | 2 | 0 | 2 |
| 109 | $2A_1 + A_{16}$ | (0) | 2 | 0 | 34 |
| | | | 4 | 2 | 18 |
| 110 | $A_2 + A_{16}$ | (0) | 6 | 3 | 10 |
| 111 | $A_1 + A_{17}$ | (0) | 4 | 2 | 10 |
| | | | $\mathbb{Z}/(3)$ | 2 | 0 |
| 112 | A_{18} | (0) | 2 | 1 | 10 |
| 113 | $2A_4 + 2D_5$ | (0) | 20 | 0 | 20 |
| 114 | $A_3 + 2A_5 + D_5$ | $\mathbb{Z}/(2)$ | 12 | 0 | 12 |
| 115 | $2A_4 + A_5 + D_5$ | (0) | 20 | 0 | 30 |
| 116 | $A_1 + A_3 + A_4 + A_5 + D_5$ | $\mathbb{Z}/(2)$ | 12 | 0 | 20 |
| 117 | $A_1 + 2A_6 + D_5$ | (0) | 14 | 0 | 28 |
| 118 | $2A_2 + A_3 + A_6 + D_5$ | (0) | 12 | 0 | 84 |
| 119 | $A_1 + A_2 + A_4 + A_6 + D_5$ | (0) | 20 | 0 | 42 |

Table 2. List of extremal elliptic $K3$ surfaces

| No | Σ | MW | a | b | c |
|-----|-------------------------------|--|-----|-----|-----|
| 120 | $A_2 + A_5 + A_6 + D_5$ | (0) | 6 | 0 | 84 |
| | | | 12 | 0 | 42 |
| 121 | $A_1 + A_7 + 2D_5$ | $\mathbb{Z}/(4)$ | 2 | 0 | 8 |
| 122 | $A_1 + A_2 + A_3 + A_7 + D_5$ | $\mathbb{Z}/(4)$ | 6 | 0 | 8 |
| 123 | $2A_1 + A_4 + A_7 + D_5$ | $\mathbb{Z}/(2)$ | 8 | 0 | 20 |
| 124 | $A_8 + 2D_5$ | (0) | 8 | 4 | 20 |
| 125 | $A_1 + A_4 + A_8 + D_5$ | (0) | 2 | 0 | 180 |
| | | | 18 | 0 | 20 |
| 126 | $A_5 + A_8 + D_5$ | (0) | 12 | 0 | 18 |
| 127 | $2A_2 + A_9 + D_5$ | (0) | 6 | 0 | 60 |
| 128 | $2A_1 + A_2 + A_9 + D_5$ | $\mathbb{Z}/(2)$ | 2 | 0 | 60 |
| 129 | $A_1 + A_3 + A_9 + D_5$ | $\mathbb{Z}/(2)$ | 8 | 4 | 12 |
| 130 | $A_4 + A_9 + D_5$ | (0) | 10 | 0 | 20 |
| 131 | $A_1 + A_2 + A_{10} + D_5$ | (0) | 14 | 4 | 20 |
| 132 | $2A_1 + A_{11} + D_5$ | $\mathbb{Z}/(4)$ | 2 | 0 | 6 |
| 133 | $A_2 + A_{11} + D_5$ | $\mathbb{Z}/(2)$ | 6 | 0 | 6 |
| 134 | $A_1 + A_{12} + D_5$ | (0) | 2 | 0 | 52 |
| | | | 6 | 2 | 18 |
| 135 | $A_{13} + D_5$ | (0) | 6 | 2 | 10 |
| 136 | $3D_6$ | $\mathbb{Z}/(2) \times \mathbb{Z}/(2)$ | 2 | 0 | 2 |
| 137 | $2A_3 + 2D_6$ | $\mathbb{Z}/(2) \times \mathbb{Z}/(2)$ | 4 | 0 | 4 |
| 138 | $2A_2 + 2A_4 + D_6$ | (0) | 30 | 0 | 30 |
| 139 | $2A_1 + 2A_5 + D_6$ | $\mathbb{Z}/(2) \times \mathbb{Z}/(2)$ | 6 | 0 | 6 |
| 140 | $A_1 + 2A_3 + A_5 + D_6$ | $\mathbb{Z}/(2) \times \mathbb{Z}/(2)$ | 4 | 0 | 12 |
| 141 | $A_3 + A_4 + A_5 + D_6$ | $\mathbb{Z}/(2)$ | 4 | 0 | 30 |
| 142 | $2A_6 + D_6$ | (0) | 14 | 0 | 14 |
| 143 | $A_2 + A_4 + A_6 + D_6$ | (0) | 6 | 0 | 70 |
| 144 | $A_1 + 2A_2 + A_7 + D_6$ | $\mathbb{Z}/(2)$ | 6 | 0 | 24 |
| 145 | $A_2 + A_3 + A_7 + D_6$ | $\mathbb{Z}/(2)$ | 4 | 0 | 24 |
| 146 | $A_1 + A_4 + A_7 + D_6$ | $\mathbb{Z}/(2)$ | 6 | 2 | 14 |
| 147 | $A_4 + A_8 + D_6$ | (0) | 4 | 2 | 46 |
| 148 | $A_1 + A_2 + A_9 + D_6$ | $\mathbb{Z}/(2)$ | 6 | 0 | 10 |
| | | $\mathbb{Z}/(2)$ | 4 | 2 | 16 |
| 149 | $A_3 + A_9 + D_6$ | $\mathbb{Z}/(2)$ | 4 | 0 | 10 |
| 150 | $A_2 + A_{10} + D_6$ | (0) | 6 | 0 | 22 |
| 151 | $A_1 + A_{11} + D_6$ | $\mathbb{Z}/(2)$ | 4 | 0 | 6 |

Table 2. List of extremal elliptic $K3$ surfaces

| No | Σ | MW | a | b | c |
|-----|--------------------------|--|-----|-----|-----|
| 152 | $A_{12} + D_6$ | (0) | 4 | 2 | 14 |
| 153 | $A_2 + A_5 + D_5 + D_6$ | $\mathbb{Z}/(2)$ | 6 | 0 | 12 |
| 154 | $A_7 + D_5 + D_6$ | $\mathbb{Z}/(2)$ | 4 | 0 | 8 |
| 155 | $2A_2 + 2D_7$ | (0) | 12 | 0 | 12 |
| 156 | $A_2 + 3A_3 + D_7$ | $\mathbb{Z}/(4)$ | 8 | 4 | 8 |
| 157 | $A_1 + A_2 + 2A_4 + D_7$ | (0) | 10 | 0 | 60 |
| 158 | $A_2 + A_3 + A_6 + D_7$ | (0) | 8 | 4 | 44 |
| 159 | $A_1 + A_4 + A_6 + D_7$ | (0) | 4 | 0 | 70 |
| 160 | $A_5 + A_6 + D_7$ | (0) | 2 | 0 | 84 |
| 161 | $2A_1 + A_2 + A_7 + D_7$ | $\mathbb{Z}/(2)$ | 4 | 0 | 24 |
| 162 | $A_1 + A_3 + A_7 + D_7$ | $\mathbb{Z}/(4)$ | 2 | 0 | 8 |
| 163 | $2A_1 + A_9 + D_7$ | $\mathbb{Z}/(2)$ | 4 | 0 | 10 |
| 164 | $A_2 + A_9 + D_7$ | (0) | 2 | 0 | 60 |
| 165 | $A_1 + A_{10} + D_7$ | (0) | 4 | 0 | 22 |
| 166 | $A_{11} + D_7$ | $\mathbb{Z}/(4)$ | 2 | 1 | 2 |
| 167 | $A_1 + A_5 + D_5 + D_7$ | $\mathbb{Z}/(2)$ | 4 | 0 | 12 |
| 168 | $A_5 + D_6 + D_7$ | $\mathbb{Z}/(2)$ | 2 | 0 | 12 |
| 169 | $2A_1 + 2D_8$ | $\mathbb{Z}/(2) \times \mathbb{Z}/(2)$ | 2 | 0 | 2 |
| 170 | $2A_2 + 2A_3 + D_8$ | $\mathbb{Z}/(2)$ | 12 | 0 | 12 |
| 171 | $2A_5 + D_8$ | $\mathbb{Z}/(2)$ | 6 | 0 | 6 |
| 172 | $2A_1 + A_3 + A_5 + D_8$ | $\mathbb{Z}/(2) \times \mathbb{Z}/(2)$ | 2 | 0 | 12 |
| 173 | $A_1 + A_4 + A_5 + D_8$ | $\mathbb{Z}/(2)$ | 2 | 0 | 30 |
| 174 | $2A_2 + A_6 + D_8$ | (0) | 12 | 6 | 24 |
| 175 | $A_1 + A_2 + A_7 + D_8$ | $\mathbb{Z}/(2)$ | 2 | 0 | 24 |
| 176 | $A_1 + A_9 + D_8$ | $\mathbb{Z}/(2)$ | 2 | 0 | 10 |
| 177 | $2D_5 + D_8$ | $\mathbb{Z}/(2)$ | 4 | 0 | 4 |
| 178 | $A_1 + A_3 + D_6 + D_8$ | $\mathbb{Z}/(2) \times \mathbb{Z}/(2)$ | 2 | 0 | 4 |
| 179 | $2D_9$ | (0) | 4 | 0 | 4 |
| 180 | $A_1 + 2A_2 + A_4 + D_9$ | (0) | 12 | 0 | 30 |
| 181 | $A_1 + A_3 + A_5 + D_9$ | $\mathbb{Z}/(2)$ | 4 | 0 | 12 |
| 182 | $A_4 + A_5 + D_9$ | (0) | 4 | 0 | 30 |
| 183 | $A_1 + A_2 + A_6 + D_9$ | (0) | 4 | 0 | 42 |
| 184 | $2A_1 + A_7 + D_9$ | $\mathbb{Z}/(2)$ | 4 | 0 | 8 |
| 185 | $A_1 + A_8 + D_9$ | (0) | 4 | 0 | 18 |
| 186 | $A_9 + D_9$ | (0) | 4 | 0 | 10 |
| 187 | $A_4 + D_5 + D_9$ | (0) | 4 | 0 | 20 |

Table 2. List of extremal elliptic $K3$ surfaces

| No | Σ | MW | a | b | c |
|-----|----------------------------|--|-----|-----|-----|
| 188 | $2A_1 + 2A_3 + D_{10}$ | $\mathbb{Z}/(2) \times \mathbb{Z}/(2)$ | 4 | 0 | 4 |
| 189 | $2A_4 + D_{10}$ | (0) | 10 | 0 | 10 |
| 190 | $A_1 + A_3 + A_4 + D_{10}$ | $\mathbb{Z}/(2)$ | 2 | 0 | 20 |
| 191 | $3A_1 + A_5 + D_{10}$ | $\mathbb{Z}/(2) \times \mathbb{Z}/(2)$ | 4 | 2 | 4 |
| 192 | $A_3 + A_5 + D_{10}$ | $\mathbb{Z}/(2)$ | 2 | 0 | 12 |
| 193 | $A_2 + A_6 + D_{10}$ | (0) | 2 | 0 | 42 |
| 194 | $A_8 + D_{10}$ | (0) | 2 | 0 | 18 |
| 195 | $A_1 + A_2 + D_5 + D_{10}$ | $\mathbb{Z}/(2)$ | 4 | 0 | 6 |
| 196 | $A_2 + D_6 + D_{10}$ | $\mathbb{Z}/(2)$ | 2 | 0 | 6 |
| 197 | $A_1 + D_7 + D_{10}$ | $\mathbb{Z}/(2)$ | 2 | 0 | 4 |
| 198 | $2A_2 + A_3 + D_{11}$ | (0) | 12 | 0 | 12 |
| 199 | $A_1 + A_2 + A_4 + D_{11}$ | (0) | 6 | 0 | 20 |
| 200 | $A_2 + A_5 + D_{11}$ | (0) | 6 | 0 | 12 |
| 201 | $A_1 + A_6 + D_{11}$ | (0) | 6 | 2 | 10 |
| 202 | $2A_1 + 2A_2 + D_{12}$ | $\mathbb{Z}/(2)$ | 6 | 0 | 6 |
| 203 | $A_1 + A_2 + A_3 + D_{12}$ | $\mathbb{Z}/(2)$ | 4 | 0 | 6 |
| 204 | $2A_1 + A_4 + D_{12}$ | $\mathbb{Z}/(2)$ | 4 | 2 | 6 |
| 205 | $A_1 + D_5 + D_{12}$ | $\mathbb{Z}/(2)$ | 2 | 0 | 4 |
| 206 | $D_6 + D_{12}$ | $\mathbb{Z}/(2)$ | 2 | 0 | 2 |
| 207 | $A_1 + A_4 + D_{13}$ | (0) | 2 | 0 | 20 |
| 208 | $A_5 + D_{13}$ | (0) | 2 | 0 | 12 |
| 209 | $D_5 + D_{13}$ | (0) | 4 | 0 | 4 |
| 210 | $2A_2 + D_{14}$ | (0) | 6 | 0 | 6 |
| 211 | $2A_1 + A_2 + D_{14}$ | $\mathbb{Z}/(2)$ | 2 | 0 | 6 |
| 212 | $A_1 + A_3 + D_{14}$ | $\mathbb{Z}/(2)$ | 2 | 0 | 4 |
| 213 | $A_4 + D_{14}$ | (0) | 4 | 2 | 6 |
| 214 | $A_1 + A_2 + D_{15}$ | (0) | 4 | 0 | 6 |
| 215 | $2A_1 + D_{16}$ | $\mathbb{Z}/(2)$ | 2 | 0 | 2 |
| 216 | $A_2 + D_{16}$ | $\mathbb{Z}/(2)$ | 2 | 1 | 2 |
| 217 | $A_1 + D_{17}$ | (0) | 2 | 0 | 4 |
| 218 | D_{18} | (0) | 2 | 0 | 2 |
| 219 | $3E_6$ | $\mathbb{Z}/(3)$ | 2 | 1 | 2 |
| 220 | $2A_3 + 2E_6$ | (0) | 12 | 0 | 12 |
| 221 | $A_1 + A_3 + 2A_4 + E_6$ | (0) | 20 | 0 | 30 |
| 222 | $A_1 + A_5 + 2E_6$ | $\mathbb{Z}/(3)$ | 2 | 0 | 6 |
| 223 | $A_2 + 2A_5 + E_6$ | $\mathbb{Z}/(3)$ | 6 | 0 | 6 |

Table 2. List of extremal elliptic $K3$ surfaces

| No | Σ | MW | a | b | c |
|-----|-------------------------------|------------------|-----|-----|-----|
| 224 | $2A_2 + A_3 + A_5 + E_6$ | $\mathbb{Z}/(3)$ | 6 | 0 | 12 |
| 225 | $A_3 + A_4 + A_5 + E_6$ | (0) | 12 | 0 | 30 |
| 226 | $A_6 + 2E_6$ | (0) | 6 | 3 | 12 |
| 227 | $A_1 + A_2 + A_3 + A_6 + E_6$ | (0) | 6 | 0 | 84 |
| | | | 12 | 0 | 42 |
| 228 | $2A_1 + A_4 + A_6 + E_6$ | (0) | 20 | 10 | 26 |
| 229 | $A_2 + A_4 + A_6 + E_6$ | (0) | 18 | 3 | 18 |
| 230 | $A_1 + A_5 + A_6 + E_6$ | (0) | 6 | 0 | 42 |
| 231 | $A_1 + A_4 + A_7 + E_6$ | (0) | 2 | 0 | 120 |
| 232 | $A_5 + A_7 + E_6$ | (0) | 6 | 0 | 24 |
| 233 | $2A_2 + A_8 + E_6$ | $\mathbb{Z}/(3)$ | 6 | 3 | 6 |
| 234 | $2A_1 + A_2 + A_8 + E_6$ | $\mathbb{Z}/(3)$ | 2 | 0 | 18 |
| 235 | $A_1 + A_3 + A_8 + E_6$ | (0) | 12 | 0 | 18 |
| 236 | $A_4 + A_8 + E_6$ | (0) | 12 | 3 | 12 |
| 237 | $A_1 + A_2 + A_9 + E_6$ | (0) | 12 | 6 | 18 |
| 238 | $A_3 + A_9 + E_6$ | (0) | 10 | 0 | 12 |
| 239 | $2A_1 + A_{10} + E_6$ | (0) | 2 | 0 | 66 |
| 240 | $A_2 + A_{10} + E_6$ | (0) | 6 | 3 | 18 |
| 241 | $A_1 + A_{11} + E_6$ | (0) | 6 | 0 | 12 |
| | | $\mathbb{Z}/(3)$ | 2 | 0 | 4 |
| 242 | $A_{12} + E_6$ | (0) | 4 | 1 | 10 |
| 243 | $A_3 + A_4 + D_5 + E_6$ | (0) | 12 | 0 | 20 |
| 244 | $A_1 + A_6 + D_5 + E_6$ | (0) | 2 | 0 | 84 |
| 245 | $A_7 + D_5 + E_6$ | (0) | 8 | 0 | 12 |
| 246 | $D_6 + 2E_6$ | (0) | 6 | 0 | 6 |
| 247 | $A_2 + A_4 + D_6 + E_6$ | (0) | 6 | 0 | 30 |
| 248 | $A_6 + D_6 + E_6$ | (0) | 4 | 2 | 22 |
| 249 | $A_1 + A_4 + D_7 + E_6$ | (0) | 4 | 0 | 30 |
| 250 | $D_5 + D_7 + E_6$ | (0) | 4 | 0 | 12 |
| 251 | $A_4 + D_8 + E_6$ | (0) | 8 | 2 | 8 |
| 252 | $A_1 + A_2 + D_9 + E_6$ | (0) | 6 | 0 | 12 |
| 253 | $A_3 + D_9 + E_6$ | (0) | 4 | 0 | 12 |
| 254 | $A_1 + D_{11} + E_6$ | (0) | 2 | 0 | 12 |
| 255 | $D_{12} + E_6$ | (0) | 4 | 2 | 4 |
| 256 | $2A_2 + 2E_7$ | (0) | 6 | 0 | 6 |
| 257 | $A_1 + A_3 + 2E_7$ | $\mathbb{Z}/(2)$ | 2 | 0 | 4 |

Table 2. List of extremal elliptic $K3$ surfaces

| No | Σ | MW | a | b | c |
|-----|-------------------------------|------------------|-----|-----|-----|
| 258 | $A_4 + 2E_7$ | (0) | 4 | 2 | 6 |
| 259 | $A_1 + 2A_3 + A_4 + E_7$ | $\mathbb{Z}/(2)$ | 4 | 0 | 20 |
| 260 | $2A_2 + A_3 + A_4 + E_7$ | (0) | 12 | 0 | 30 |
| 261 | $2A_3 + A_5 + E_7$ | $\mathbb{Z}/(2)$ | 4 | 0 | 12 |
| 262 | $A_1 + A_2 + A_3 + A_5 + E_7$ | $\mathbb{Z}/(2)$ | 6 | 0 | 12 |
| 263 | $2A_1 + A_4 + A_5 + E_7$ | $\mathbb{Z}/(2)$ | 8 | 2 | 8 |
| 264 | $A_2 + A_4 + A_5 + E_7$ | (0) | 6 | 0 | 30 |
| 265 | $A_1 + 2A_2 + A_6 + E_7$ | (0) | 6 | 0 | 42 |
| 266 | $A_2 + A_3 + A_6 + E_7$ | (0) | 4 | 0 | 42 |
| 267 | $A_1 + A_4 + A_6 + E_7$ | (0) | 2 | 0 | 70 |
| | | | 8 | 2 | 18 |
| 268 | $A_5 + A_6 + E_7$ | (0) | 4 | 2 | 22 |
| 269 | $2A_2 + A_7 + E_7$ | (0) | 6 | 0 | 24 |
| 270 | $2A_1 + A_2 + A_7 + E_7$ | $\mathbb{Z}/(2)$ | 2 | 0 | 24 |
| 271 | $A_1 + A_3 + A_7 + E_7$ | $\mathbb{Z}/(2)$ | 4 | 0 | 8 |
| 272 | $A_4 + A_7 + E_7$ | (0) | 6 | 2 | 14 |
| 273 | $A_1 + A_2 + A_8 + E_7$ | (0) | 6 | 0 | 18 |
| 274 | $A_3 + A_8 + E_7$ | (0) | 4 | 0 | 18 |
| 275 | $2A_1 + A_9 + E_7$ | $\mathbb{Z}/(2)$ | 2 | 0 | 10 |
| 276 | $A_2 + A_9 + E_7$ | (0) | 6 | 0 | 10 |
| | | $\mathbb{Z}/(2)$ | 4 | 1 | 4 |
| 277 | $A_1 + A_{10} + E_7$ | (0) | 2 | 0 | 22 |
| | | | 6 | 2 | 8 |
| 278 | $A_{11} + E_7$ | (0) | 4 | 0 | 6 |
| 279 | $D_4 + 2E_7$ | $\mathbb{Z}/(2)$ | 2 | 0 | 2 |
| 280 | $A_2 + A_4 + D_5 + E_7$ | (0) | 6 | 0 | 20 |
| 281 | $A_1 + A_5 + D_5 + E_7$ | $\mathbb{Z}/(2)$ | 2 | 0 | 12 |
| 282 | $A_6 + D_5 + E_7$ | (0) | 6 | 2 | 10 |
| 283 | $A_2 + A_3 + D_6 + E_7$ | $\mathbb{Z}/(2)$ | 4 | 0 | 6 |
| 284 | $A_5 + D_6 + E_7$ | $\mathbb{Z}/(2)$ | 4 | 2 | 4 |
| 285 | $D_5 + D_6 + E_7$ | $\mathbb{Z}/(2)$ | 2 | 0 | 4 |
| 286 | $A_1 + A_3 + D_7 + E_7$ | $\mathbb{Z}/(2)$ | 4 | 0 | 4 |
| 287 | $A_4 + D_7 + E_7$ | (0) | 2 | 0 | 20 |
| 288 | $A_1 + A_2 + D_8 + E_7$ | $\mathbb{Z}/(2)$ | 2 | 0 | 6 |
| 289 | $A_2 + D_9 + E_7$ | (0) | 4 | 0 | 6 |
| 290 | $A_1 + D_{10} + E_7$ | $\mathbb{Z}/(2)$ | 2 | 0 | 2 |

Table 2. List of extremal elliptic $K3$ surfaces

| No | Σ | MW | a | b | c |
|-----|-------------------------------|------|-----|-----|-----|
| 291 | $D_{11} + E_7$ | (0) | 2 | 0 | 4 |
| 292 | $A_2 + A_3 + E_6 + E_7$ | (0) | 6 | 0 | 12 |
| 293 | $A_1 + A_4 + E_6 + E_7$ | (0) | 2 | 0 | 30 |
| 294 | $A_5 + E_6 + E_7$ | (0) | 6 | 0 | 6 |
| 295 | $D_5 + E_6 + E_7$ | (0) | 2 | 0 | 12 |
| 296 | $2A_1 + 2E_8$ | (0) | 2 | 0 | 2 |
| 297 | $A_2 + 2E_8$ | (0) | 2 | 1 | 2 |
| 298 | $2A_2 + 2A_3 + E_8$ | (0) | 12 | 0 | 12 |
| 299 | $2A_1 + 2A_4 + E_8$ | (0) | 10 | 0 | 10 |
| 300 | $A_1 + A_2 + A_3 + A_4 + E_8$ | (0) | 6 | 0 | 20 |
| 301 | $2A_5 + E_8$ | (0) | 6 | 0 | 6 |
| 302 | $A_2 + A_3 + A_5 + E_8$ | (0) | 6 | 0 | 12 |
| 303 | $A_1 + A_4 + A_5 + E_8$ | (0) | 2 | 0 | 30 |
| 304 | $2A_2 + A_6 + E_8$ | (0) | 6 | 3 | 12 |
| 305 | $2A_1 + A_2 + A_6 + E_8$ | (0) | 2 | 0 | 42 |
| 306 | $A_1 + A_3 + A_6 + E_8$ | (0) | 6 | 2 | 10 |
| 307 | $A_4 + A_6 + E_8$ | (0) | 2 | 1 | 18 |
| 308 | $A_1 + A_2 + A_7 + E_8$ | (0) | 2 | 0 | 24 |
| 309 | $2A_1 + A_8 + E_8$ | (0) | 2 | 0 | 18 |
| 310 | $A_2 + A_8 + E_8$ | (0) | 6 | 3 | 6 |
| 311 | $A_1 + A_9 + E_8$ | (0) | 2 | 0 | 10 |
| 312 | $A_{10} + E_8$ | (0) | 2 | 1 | 6 |
| 313 | $2D_5 + E_8$ | (0) | 4 | 0 | 4 |
| 314 | $A_1 + A_4 + D_5 + E_8$ | (0) | 2 | 0 | 20 |
| 315 | $A_5 + D_5 + E_8$ | (0) | 2 | 0 | 12 |
| 316 | $2A_2 + D_6 + E_8$ | (0) | 6 | 0 | 6 |
| 317 | $A_4 + D_6 + E_8$ | (0) | 4 | 2 | 6 |
| 318 | $A_1 + A_2 + D_7 + E_8$ | (0) | 4 | 0 | 6 |
| 319 | $A_1 + D_9 + E_8$ | (0) | 2 | 0 | 4 |
| 320 | $D_{10} + E_8$ | (0) | 2 | 0 | 2 |
| 321 | $A_1 + A_3 + E_6 + E_8$ | (0) | 2 | 0 | 12 |
| 322 | $A_4 + E_6 + E_8$ | (0) | 2 | 1 | 8 |
| 323 | $D_4 + E_6 + E_8$ | (0) | 4 | 2 | 4 |
| 324 | $A_1 + A_2 + E_7 + E_8$ | (0) | 2 | 0 | 6 |
| 325 | $A_3 + E_7 + E_8$ | (0) | 2 | 0 | 4 |

REFERENCES

1. E. Artal-Bartolo, H. Tokunaga and D. Q. Zhang. *Miranda-Persson's problem on extremal elliptic $K3$ surfaces*. preprint. <http://xxx.lanl.gov/list/math.AG,9809065>.
2. N. Bourbaki. *Éléments de mathématique. Groupes et algèbres de Lie. Chapitre IV-VI*. Hermann, Paris, 1968.
3. J. H. Conway and N. J. A. Sloane. *Sphere packings, lattices and groups*. Second edition. Grundlehren der Mathematischen Wissenschaften, **290**, Springer, New York, 1993.
4. A. Fujiki. *Finite automorphism groups of complex tori of dimension two*. Publ. Res. Inst. Math. Sci. **24** (1988), no. 1, 1–97.
5. S. Kondō. *Automorphisms of algebraic $K3$ surfaces which act trivially on Picard groups*. J. Math. Soc. Japan **44** (1992), no. 1, 75–98.
6. S. Kondō. *Niemeyer lattices, Mathieu groups, and finite groups of symplectic automorphisms of $K3$ surfaces*. With an appendix by Shigeru Mukai. Duke Math. J. **92** (1998), no. 3, 593–603.
7. R. Miranda and U. Persson. *Mordell-Weil groups of extremal elliptic $K3$ surfaces*. Problems in the theory of surfaces and their classification (Cortona, 1988), Sympos. Math., XXXII, Academic Press, London, 1991, pp. 167–192.
8. D. R. Morrison. *On $K3$ surfaces with large Picard number*. Invent. Math. **75** (1984), no. 1, 105–121.
9. S. Mukai. *Finite groups of automorphisms of $K3$ surfaces and the Mathieu group*. Invent. Math. **94** (1988), no. 1, 183–221.
10. V. V. Nikulin. *Finite automorphism groups of Kähler $K3$ surfaces*. Trans. Moscow Math. Soc. (1980), Issue 2, pp. 71–135.
11. V. V. Nikulin. *Integer symmetric bilinear forms and some of their applications*. Math. USSR Izvestija **14** (1980), no. 1, 103–167.
12. K. Nishiyama. *The Jacobian fibrations on some $K3$ surfaces and their Mordell-Weil groups*. Japan. J. Math. (N.S.) **22** (1996), no. 2, 293–347.
13. M. V. Nori. *Zariski's conjecture and related problems*. Ann. Sci. École Norm. Sup. (4) **16** (1983), no. 2, 305–344.
14. I. Piatetskii-Shapiro and I. R. Shafarevich. *A Torelli theorem for algebraic surfaces of type $K3$* . Math. USSR Izv. **35** (1971), 530–572.
15. J.-P. Serre. *A course in arithmetic*. Graduate Texts in Mathematics, **7**, Springer, New York, 1973.
16. T. Shioda and H. Inose. *On singular $K3$ surfaces*. Complex analysis and algebraic geometry, Iwanami Shoten, Tokyo, 1977, pp. 119–136.
17. A. N. Todorov. *Applications of the Kähler-Einstein-Calabi-Yau metric to moduli of $K3$ surfaces*. Invent. Math. **61** (1980), no. 3, 251–265.
18. G. Xiao. *Galois covers between $K3$ surfaces*. Ann. Inst. Fourier (Grenoble) **46** (1996), no. 1, 73–88.
19. Q. Ye. *On extremal elliptic $K3$ surfaces*. preprint. <http://xxx.lanl.gov/abs/math.AG,9901081>

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, SAPPORO, JAPAN 061-0081
E-mail address: shimada@math.sci.hokudai.ac.jp

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, LOWER KENTRIDGE ROAD, SINGAPORE 119260
E-mail address: matzdzq@math.nus.edu.sg