

The intersection of two real flag manifolds in a complex flag manifold

Takashi Sakai (Tokyo Metropolitan University / OCAMI)

joint work with Osamu Ikawa, Hiroshi Iriyeh,
Takayuki Okuda, and Hiroyuki Tasaki

December 13, 2016

The First Japan-Taiwan Joint Conference on Differential Geometry &
the 8th TIMS-OCAMI-WASEDA Joint International Workshop
on Differential Geometry and Geometric Analysis
at Waseda University

(M, J, ω) : homogeneous Kähler manifold

L_1, L_2 : real forms of M

i.e. $\exists \sigma_i$: anti-holomorphic involutive isometry of M ($i = 1, 2$)

s.t. $L_i = \text{Fix}(\sigma_i, M)_0$


totally geodesic Lagrangian submanifold

Problems

- 1 Is the intersection $L_1 \cap L_2$ discrete?
- 2 If so, count the intersection number $\#(L_1 \cap L_2)$, and describe the geometric meaning of $\#(L_1 \cap L_2)$.
Moreover, study the structure of the intersection $L_1 \cap L_2$.
- 3 Using the antipodal structure of the intersection, study the Floer homology of L_1 and L_2 .

Problems

- 1 Is the intersection $L_1 \cap L_2$ discrete?
- 2 If so, count the intersection number $\#(L_1 \cap L_2)$, and describe the geometric meaning of $\#(L_1 \cap L_2)$.
Moreover, study the structure of the intersection $L_1 \cap L_2$.




$$M = \mathbb{C}P^1$$

$$L_1 = \mathbb{R}P^1,$$

Problems

- 1 Is the intersection $L_1 \cap L_2$ discrete?
- 2 If so, count the intersection number $\#(L_1 \cap L_2)$, and describe the geometric meaning of $\#(L_1 \cap L_2)$.

Moreover, study the structure of the intersection $L_1 \cap L_2$.



$$M = \mathbb{C}P^1$$

$$L_1 = \mathbb{R}P^1, \quad L_2 \cong \mathbb{R}P^1$$


$$\#(L_1 \cap L_2) = 2 = \dim H_*(L_1; \mathbb{Z}_2)$$

$L_1 \cap L_2$: antipodal points

Problems

- 1 Is the intersection $L_1 \cap L_2$ discrete?
- 2 If so, count the intersection number $\#(L_1 \cap L_2)$, and describe the geometric meaning of $\#(L_1 \cap L_2)$.

Moreover, study the structure of the intersection $L_1 \cap L_2$.



$$M = \mathbb{C}P^1$$

$$L_1 = \mathbb{R}P^1, \quad L_2 \cong \mathbb{R}P^1$$

$$\#(L_1 \cap L_2) = 2 = \dim H_*(L_1; \mathbb{Z}_2)$$

$L_1 \cap L_2$: antipodal points

$$\#(L_1 \cap \phi L_2) \geq 2 \quad \forall \phi \in \text{Ham}(M, \omega)$$

Theorem (Tanaka-Tasaki 2012)

M : Hermitian symmetric space of compact type

$L_1, L_2 \subset M$: real forms, $L_1 \pitchfork L_2$

$\implies L_1 \cap L_2$ is an *antipodal set* of L_1 and L_2 .

In addition, if L_1 and L_2 are congruent to each other,

$\implies L_1 \cap L_2$ is a *great antipodal set* of L_1 and L_2 .

Theorem (Ikawa-Tanaka-Tasaki 2015)

A necessary and sufficient condition for two real forms in a compact Hermitian symmetric space to intersect transversally is given in terms of *symmetric triad* $(\tilde{\Sigma}, \Sigma, W)$.

Antipodal sets of a compact symmetric space (1/2)

M : compact Riemannian symmetric space

s_x : geodesic symmetry at $x \in M$

Definition (Chen-Nagano 1988)

- 1 $\mathcal{A} \subset M$: **antipodal set** $\stackrel{\text{def}}{\iff} s_x(y) = y$ for all $x, y \in \mathcal{A}$
- 2 $\#_2 M := \max\{\#\mathcal{A} \mid \mathcal{A} \subset M : \text{antipodal set}\}$ **2-number**
- 3 $\mathcal{A} \subset M$: **great antipodal set** $\stackrel{\text{def}}{\iff} \#\mathcal{A} = \#_2 M$

Theorem (Takeuchi 1989)

M : *symmetric R-space* $\implies \#_2 M = \dim H_*(M; \mathbb{Z}_2)$

Antipodal sets of a compact symmetric space (2/2)

Example

$$\mathbb{R}P^n \subset \mathbb{C}P^n$$

$$\mathcal{A} := \{\mathbb{R}e_1, \dots, \mathbb{R}e_{n+1}\} \subset \mathbb{R}P^n \quad \text{great antipodal set}$$

For $u \in U(n+1)$, $\mathbb{R}P^n \pitchfork u\mathbb{R}P^n$ in $\mathbb{C}P^n$

$$\mathbb{R}P^n \cap u\mathbb{R}P^n \cong \{\mathbb{C}e_1, \dots, \mathbb{C}e_{n+1}\} \subset \mathbb{C}P^n$$

$$\#(\mathbb{R}P^n \cap u\mathbb{R}P^n) = n + 1 = \#_2 \mathbb{R}P^n = \dim H_*(\mathbb{R}P^n; \mathbb{Z}_2)$$

Aim of our work

Generalizing the results on Hermitian symmetric spaces, study the intersection of two real forms in a complex flag manifold.

Complex flag manifolds

G : compact connected semisimple Lie group

$x_0 (\neq 0) \in \mathfrak{g}$

$$\begin{aligned} M &:= \text{Ad}(G)x_0 \subset \mathfrak{g} && : \text{complex flag manifold} \\ &\cong G/G_{x_0} \cong G^{\mathbb{C}}/P^{\mathbb{C}} \end{aligned}$$

$$G_{x_0} := \{g \in G \mid \text{Ad}(g)x_0 = x_0\}$$

ω : Kirillov-Kostant-Souriau symplectic form on M defined by

$$\omega(X_x^*, Y_x^*) := \langle [X, Y], x \rangle \quad (x \in M, X, Y \in \mathfrak{g})$$

J : G -invariant complex structure on M compatible with ω

$(\cdot, \cdot) := \omega(\cdot, J\cdot)$: G -invariant Kähler metric

Antipodal set of a complex flag manifold (1/2)

For $x \in M$ and $g \in Z(G_{x_0})$, define $s_{x,g} : M \rightarrow M$ by

$$s_{x,g}(y) := \text{Ad}(g_x g g_x^{-1})y \quad (y \in M),$$

where $g_x \in G$ satisfying $\text{Ad}(g_x)x_0 = x$.

$$\text{Fix}(s_x, M) := \{y \in M \mid s_{x,g}(y) = y \ (\forall g \in Z(G_{x_0}))\}$$

Definition

$\mathcal{A} \subset M$: **antipodal set** \iff $y \in \text{Fix}(s_x, M)$ for all $x, y \in \mathcal{A}$

Antipodal set of a complex flag manifold (2/2)

Proposition 1

For any $x \in M$,

$$\text{Fix}(s_x, M) = \{y \in M \mid [x, y] = 0\}.$$

Theorem 1 (Iriyeh-S.-Tasaki)

$\mathcal{A} \subset M$: maximal antipodal set

$\implies \exists \mathfrak{t} \subset \mathfrak{g}$: maximal abelian subalgebra s.t.

$$\mathcal{A} = M \cap \mathfrak{t}.$$

Hence \mathcal{A} is an orbit of the Weyl group of \mathfrak{g} with respect to \mathfrak{t} .

Maximal antipodal sets of M are congruent to each other by G .

Real flag manifolds in a complex flag manifold

(G, K) : symmetric pair of compact type

θ : involution of G s.t. $\text{Fix}(\theta, G)_0 \subset K \subset \text{Fix}(\theta, G)$

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$$

$x_0 (\neq 0) \in \mathfrak{p}$

$L := \text{Ad}(K)x_0 \subset \mathfrak{p}$: **real flag manifold, R -space**

$\cap \quad \cap \quad \cap$

$M := \text{Ad}(G)x_0 \subset \mathfrak{g}$: **complex flag manifold, C -space**

$$\cong G/G_{x_0} \cong G^{\mathbb{C}}/P^{\mathbb{C}}$$

$\mathfrak{g}' := \mathfrak{k} + \sqrt{-1}\mathfrak{p}$ non-compact real form of $\mathfrak{g}^{\mathbb{C}}$

$$L = M \cap \mathfrak{p} \cong K/K_{x_0} \cong G'/(G' \cap P^{\mathbb{C}})$$

The intersection of real flag manifolds

$(G, K_1), (G, K_2)$: symmetric pairs of compact type

θ_1, θ_2 : involutions of G

$$\mathfrak{g} = \mathfrak{k}_1 + \mathfrak{p}_1 = \mathfrak{k}_2 + \mathfrak{p}_2,$$

$$x_0 (\neq 0) \in \mathfrak{p}_1 \cap \mathfrak{p}_2$$

$$L_1 := \text{Ad}(K_1)x_0, \quad L_2 := \text{Ad}(K_2)x_0 \subset M := \text{Ad}(G)x_0$$

For $g \in G$, we consider the intersection of $L_1 \cap \text{Ad}(g)L_2$ in M .

\mathfrak{a} : maximal abelian subspace of $\mathfrak{p}_1 \cap \mathfrak{p}_2$ containing x_0

$A := \exp \mathfrak{a} \subset G$: toral subgroup

Then $G = K_1AK_2$, i.e. $g = g_1ag_2$ ($g_1 \in K_1, g_2 \in K_2, a \in A$)

$$L_1 \cap \text{Ad}(g)L_2 = L_1 \cap \text{Ad}(g_1ag_2)L_2 = \text{Ad}(g_1) \left(L_1 \cap \text{Ad}(a)L_2 \right)$$

Symmetric triads

Hereafter we assume that G is simple and $\theta_1\theta_2 = \theta_2\theta_1$.

$$\mathfrak{g} = \underbrace{(\mathfrak{k}_1 \cap \mathfrak{k}_2) + (\mathfrak{p}_1 \cap \mathfrak{p}_2)}_{\text{ad}(\mathfrak{a})\text{-inv.}} + \underbrace{(\mathfrak{k}_1 \cap \mathfrak{p}_2) + (\mathfrak{p}_1 \cap \mathfrak{k}_2)}_{\text{ad}(\mathfrak{a})\text{-inv.}}$$

$\mathfrak{a} \subset \mathfrak{p}_1 \cap \mathfrak{p}_2$: maximal abelian subspace

For $\lambda \in \mathfrak{a}$

$$\mathfrak{p}_\lambda := \{X \in \mathfrak{p}_1 \cap \mathfrak{p}_2 \mid (\text{ad}H)^2 X = -\langle \lambda, H \rangle^2 X \ (H \in \mathfrak{a})\}$$

$$V_\lambda := \{X \in \mathfrak{p}_1 \cap \mathfrak{k}_2 \mid (\text{ad}H)^2 X = -\langle \lambda, H \rangle^2 X \ (H \in \mathfrak{a})\}$$

$$\Sigma := \{\lambda \in \mathfrak{a} \setminus \{0\} \mid \mathfrak{p}_\lambda \neq \{0\}\}$$

$$W := \{\lambda \in \mathfrak{a} \setminus \{0\} \mid V_\lambda \neq \{0\}\}$$

$$\tilde{\Sigma} := \Sigma \cup W$$

$(\tilde{\Sigma}, \Sigma, W)$: **symmetric triad** with multiplicities

The structure of the intersection

$$\mathfrak{a}_{\text{reg}} := \bigcap_{\substack{\lambda \in \Sigma \\ \alpha \in W}} \left\{ H \in \mathfrak{a} \mid \langle \lambda, H \rangle \notin \pi\mathbb{Z}, \langle \alpha, H \rangle \notin \frac{\pi}{2} + \pi\mathbb{Z} \right\}$$

$W(\tilde{\Sigma})$: Weyl group of the root system $\tilde{\Sigma}$ of \mathfrak{a}

\mathfrak{a}_i : maximal abelian subspace of \mathfrak{p}_i containing \mathfrak{a} ($i = 1, 2$)

$W(R_i)$: Weyl group of the restricted root system R_i of $(\mathfrak{g}, \mathfrak{k}_i)$
w.r.t. \mathfrak{a}_i

Theorem 2 (Ikawa-Iriyeh-Okuda-S.-Tasaki)

For $a = \exp H$ ($H \in \mathfrak{a}$), the intersection $L_1 \cap \text{Ad}(a)L_2$ is discrete if and only if $H \in \mathfrak{a}_{\text{reg}}$. Moreover, if $L_1 \cap \text{Ad}(a)L_2$ is discrete, then

$$L_1 \cap \text{Ad}(a)L_2 = W(\tilde{\Sigma})x_0 = W(R_1)x_0 \cap \mathfrak{a} = W(R_2)x_0 \cap \mathfrak{a}.$$

In particular, $L_1 \cap \text{Ad}(a)L_2$ is an antipodal set of M .

Example

$$(G, K_1, K_2) = (SU(2n), SO(2n), Sp(n))$$

$$\theta_1(g) = \bar{g}, \quad \theta_2(g) = J_n \bar{g} J_n^{-1} \quad (g \in G) \quad \text{where} \quad J_n := \begin{bmatrix} O & I_n \\ -I_n & O \end{bmatrix}$$

$$\mathfrak{p}_1 \cap \mathfrak{p}_2 = \left\{ \left[\begin{array}{cc} \sqrt{-1}X & \sqrt{-1}Y \\ -\sqrt{-1}Y & \sqrt{-1}X \end{array} \right] \mid \begin{array}{l} X, Y \in M_n(\mathbb{R}), \text{ trace } X = 0 \\ {}^t X = X, {}^t Y = -Y \end{array} \right\}$$

Fix a maximal abelian subspace \mathfrak{a} in $\mathfrak{p}_1 \cap \mathfrak{p}_2$ as

$$\mathfrak{a} = \left\{ H = \begin{bmatrix} \sqrt{-1}X & O \\ O & \sqrt{-1}X \end{bmatrix} \mid \begin{array}{l} X = \text{diag}(t_1, \dots, t_n), \\ t_1, \dots, t_n \in \mathbb{R}, t_1 + \dots + t_n = 0 \end{array} \right\}$$

Then

$$\tilde{\Sigma} = \Sigma = W = \{\pm(e_i - e_j) \mid 1 \leq i < j \leq n\}$$

where $e_i - e_j \in \mathfrak{a}$ ($i \neq j$) is defined by $\langle e_i - e_j, H \rangle = t_i - t_j$.

$$x_0 = \begin{bmatrix} \sqrt{-1}X & O \\ O & \sqrt{-1}X \end{bmatrix} \in \mathfrak{a}$$

where $X = \text{diag}(x_1 I_{n_1}, \dots, x_{r+1} I_{n_{r+1}})$ and x_i are distinct real numbers satisfying $n_1 x_1 + \dots + n_{r+1} x_{r+1} = 0$.

$$L_1 = \text{Ad}(K_1)x_0 \cong F_{2n_1, \dots, 2n_r}^{\mathbb{R}}(\mathbb{R}^{2n})$$

$$L_2 = \text{Ad}(K_2)x_0 \cong F_{n_1, \dots, n_r}^{\mathbb{H}}(\mathbb{H}^n)$$

$$M = \text{Ad}(G)x_0 \cong F_{2n_1, \dots, 2n_r}^{\mathbb{C}}(\mathbb{C}^{2n})$$

$\mathbb{K} = \mathbb{R}, \mathbb{C}$ or \mathbb{H}

n, n_1, \dots, n_r satisfying $n_{r+1} := n - (n_1 + \dots + n_r) > 0$

$$F_{n_1, \dots, n_r}^{\mathbb{K}}(\mathbb{K}^n) = \left\{ (V_1, \dots, V_r) \left| \begin{array}{l} V_j \text{ is a } \mathbb{K}\text{-subspace of } \mathbb{K}^n, \\ \dim_{\mathbb{K}} V_j = n_1 + \dots + n_j, \\ V_1 \subset V_2 \subset \dots \subset V_r \subset \mathbb{K}^n \end{array} \right. \right\}$$

$$a = \exp H, \quad H = \begin{bmatrix} \sqrt{-1}Y & O \\ O & \sqrt{-1}Y \end{bmatrix} \in \mathfrak{a}$$

where $Y = \text{diag}(t_1, \dots, t_n)$ and $t_1, \dots, t_n \in \mathbb{R}$ which satisfy $t_1 + \dots + t_n = 0$. By our theorem,

$L_1 \cap \text{Ad}(a)L_2$ is discrete

$$\iff H \in \mathfrak{a}_{\text{reg}} = \left\{ H \in \mathfrak{a} \mid \langle e_i - e_j, H \rangle \notin \frac{\pi}{2}\mathbb{Z} \ (1 \leq i < j \leq n) \right\}$$

$$L_1 \cap \text{Ad}(a)L_2 = W(\tilde{\Sigma})x_0 = W(R_1)x_0 \cap \mathfrak{a} = W(R_2)x_0 \cap \mathfrak{a}$$

In this case, a maximal abelian subspace \mathfrak{a} in $\mathfrak{p}_1 \cap \mathfrak{p}_2$ is also a maximal abelian subspace in \mathfrak{p}_2 , i.e. $\mathfrak{a} = \mathfrak{a}_2$ and $\tilde{\Sigma} = R_2$.

We shall express the intersection in the flag model $F_{2n_1, \dots, 2n_r}^{\mathbb{C}}(\mathbb{C}^{2n})$.

v_1, \dots, v_{2n} : standard basis of \mathbb{C}^{2n}

$W_i := \langle v_i, v_{n+i} \rangle_{\mathbb{C}} = \langle v_i \rangle_{\mathbb{H}} \quad (1 \leq i \leq n)$

Proposition 2

For $a = \exp H \quad (H \in \mathfrak{a}_{\text{reg}})$,

$$\begin{aligned}
 & F_{2n_1, \dots, 2n_r}^{\mathbb{R}}(\mathbb{R}^{2n}) \cap aF_{n_1, \dots, n_r}^{\mathbb{H}}(\mathbb{H}^n) \\
 &= \{ (W_{i_1} \oplus \cdots \oplus W_{i_{n_1}}, W_{i_1} \oplus \cdots \oplus W_{i_{n_1+n_2}}, \dots \\
 &\quad \dots, W_{i_1} \oplus \cdots \oplus W_{i_{n_1+\dots+n_r}}) \\
 & \mid 1 \leq i_1 < \cdots < i_{n_1} \leq n, 1 \leq i_{n_1+1} < \cdots < i_{n_1+n_2} \leq n, \dots, \\
 & \quad 1 \leq i_{n_1+\dots+n_{r-1}+1} < \cdots < i_{n_1+\dots+n_r} \leq n, \\
 & \quad \#\{i_1, \dots, i_{n_1+\dots+n_r}\} = n_1 + \cdots + n_r \},
 \end{aligned}$$

which is an antipodal set of $F_{2n_1, \dots, 2n_r}^{\mathbb{C}}(\mathbb{C}^{2n})$.

Corollary 1

For $a = \exp H$ ($H \in \mathfrak{a}_{\text{reg}}$)

$$\begin{aligned} & \#(F_{2n_1, \dots, 2n_r}^{\mathbb{R}}(\mathbb{R}^{2n}) \cap gF_{n_1, \dots, n_r}^{\mathbb{H}}(\mathbb{H}^n)) \\ &= \#_I(F_{n_1, \dots, n_r}^{\mathbb{H}}(\mathbb{H}^n)) = \dim H_*(F_{n_1, \dots, n_r}^{\mathbb{H}}(\mathbb{H}^n); \mathbb{Z}_2) \\ &= \frac{n!}{n_1!n_2! \cdots n_{r+1}!} \\ &< \#_I(F_{2n_1, \dots, 2n_r}^{\mathbb{R}}(\mathbb{R}^{2n})) = \dim H_*(F_{2n_1, \dots, 2n_r}^{\mathbb{R}}(\mathbb{R}^{2n}); \mathbb{Z}_2) \\ &= \#_k(F_{2n_1, \dots, 2n_r}^{\mathbb{C}}(\mathbb{C}^{2n})) = \dim H_*(F_{2n_1, \dots, 2n_r}^{\mathbb{C}}(\mathbb{C}^{2n}); \mathbb{Z}_2) \\ &= \frac{(2n)!}{(2n_1)!(2n_2)! \cdots (2n_{r+1})!}. \end{aligned}$$

Theorem (Sánchez, Berndt-Console-Fino)

For a complex flag manifold M and a real flag manifold L ,

$$\#_k(M) = \dim H_*(M; \mathbb{Z}_2), \quad \#_I(L) = \dim H_*(L; \mathbb{Z}_2).$$

Lagrangian Floer homology

(M, ω) : closed symplectic manifold

$J = \{J_t\}_{0 \leq t \leq 1}$: family of ω -compatible almost complex structures

L_0, L_1 : closed Lagrangian submanifolds, $L_0 \pitchfork L_1$

$$CF(L_0, L_1) := \bigoplus_{p \in L_0 \cap L_1} \mathbb{Z}_2 p$$

$\partial : CF(L_0, L_1) \longrightarrow CF(L_0, L_1)$

$$\partial(p) = \sum_{q \in L_0 \cap L_1} n(p, q) \cdot q$$

$n(p, q) := \#\{\text{isolated } J\text{-holomorphic strips from } p \text{ to } q\} \pmod{2}$

$$\partial \circ \partial = 0 \quad \implies \quad HF(L_0, L_1; \mathbb{Z}_2) := \ker \partial / \text{im } \partial$$

① $HF(\phi L_0, \psi L_1; \mathbb{Z}_2) \cong HF(L_0, L_1; \mathbb{Z}_2) \quad \forall \phi, \psi \in \text{Ham}(M, \omega)$

Theorem 3 (Iriyeh-S.-Tasaki 2013)

(M, J_0, ω) : Einstein, Hermitian symmetric space of compact type

L_0, L_1 : real forms, $L_0 \pitchfork L_1$, $\Sigma_{L_0}, \Sigma_{L_1} \geq 3$

\implies

$$HF(L_0, L_1; \mathbb{Z}_2) \cong \bigoplus_{p \in L_0 \cap L_1} \mathbb{Z}_2[p]$$

- 1 If M is irreducible, then the assumptions are satisfied except for the case $\mathbb{R}P^1 \subset \mathbb{C}P^1$.

Theorem 4 (Iriyeh-S.-Tasaki 2013)

M : irreducible Hermitian symmetric space of compact type

L_0, L_1 : real forms of M , $L_0 \pitchfork L_1$

\implies

$$\textcircled{1} \quad (M, L_0, L_1) \cong (G_{2m}^{\mathbb{C}}(\mathbb{C}^{4m}), G_m^{\mathbb{H}}(\mathbb{H}^{2m}), U(2m)) \quad (m \geq 2)$$

$$HF(L_0, L_1; \mathbb{Z}_2) \cong (\mathbb{Z}_2)^{2^m}$$

where $2^m < \binom{2m}{m} = \#_2 L_0 < 2^{2m} = \#_2 L_1$

$$\textcircled{2} \quad (M, L_0, L_1) : \text{otherwise}$$

$$HF(L_0, L_1; \mathbb{Z}_2) \cong (\mathbb{Z}_2)^{\min\{\#_2 L_0, \#_2 L_1\}}$$

Corollary 2

M : irreducible Hermitian symmetric space of compact type

(L_0, L_1) : real forms of M

\implies for any $\phi \in \text{Ham}(M, \omega)$, $L_0 \pitchfork \phi L_1$

① $(M, L_0, L_1) \cong (G_{2m}^{\mathbb{C}}(\mathbb{C}^{4m}), G_m^{\mathbb{H}}(\mathbb{H}^{2m}), U(2m)) \quad (m \geq 2)$

$$\#(L_0 \cap \phi L_1) \geq 2^m$$

② (M, L_0, L_1) : otherwise

$$\#(L_0 \cap \phi L_1) \geq \min\{SB(L_0, \mathbb{Z}_2), SB(L_1, \mathbb{Z}_2)\}$$

Further problems

- 1 Study the intersection of two real flag manifolds in the case where $\theta_1\theta_2 \neq \theta_2\theta_1$.
- 2 Calculate Lagrangian Floer homologies of pairs of real flag manifolds in complex flag manifolds.
- 3 Determine Hamiltonian volume minimizing properties of all real forms in irreducible Hermitian symmetric spaces, more generally, in complex flag manifolds.

Thank you very much for your attention

Further problems

- 1 Study the intersection of two real flag manifolds in the case where $\theta_1\theta_2 \neq \theta_2\theta_1$.
- 2 Calculate Lagrangian Floer homologies of pairs of real flag manifolds in complex flag manifolds.
- 3 Determine Hamiltonian volume minimizing properties of all real forms in irreducible Hermitian symmetric spaces, more generally, in complex flag manifolds.

Thank you very much for your attention