


EoS in 2+1 flavor QCD with improved Wilson fermion

Takashi Umeda (Hiroshima Univ.)
for WHOT-QCD Collaboration




*Nonperturbative Aspects of QCD at Finite Temperature and Density,
Univ. of Tsukuba, Ibaraki, 8 - 9 Nov. 2010*

Quark Gluon Plasma in Lattice QCD



from the Phenix group web-site



<http://www.gsi.de/fair/experiments/>

Observables in Lattice QCD

- Phase diagram in (T, μ, m_{ud}, m_s)
- Transition temperature
- Equation of state ($\epsilon/T^4, p/T^4, \dots$)
- Hadronic excitations
- Transport coefficients
- Finite chemical potential
- etc...

Choice of quark actions on the lattice

Most ($T, \mu \neq 0$) studies done with staggered-type quarks

- less computational costs
- a part of chiral sym. preserved ...
 - $N_f=2+1$, almost physical quark mass, ($\mu \neq 0$)
- 4th-root trick to remove unphysical "tastes"
 - non-locality "Validity is not guaranteed"

It is important to cross-check with
theoretically sound lattice quarks like Wilson-type quarks

Our aim is to calculate
QCD EoS using **Wilson-type quarks**
with **small lattice artifacts**.


Fixed scale approach to study QCD thermodynamics

Temperature $T=1/(N_t a)$ is varied by N_t at fixed a

a : lattice spacing

N_t : lattice size in temporal direction

Temperatures in each approach



■ Advantages

- Line of Constant Physics
- $T=0$ subtraction for renorm. (spectrum study at $T=0$)
- fine "a" in whole T region

■ Disadvantages

- T resolution by integer N_t
- UV cutoff eff. at high T

T-integration method to calculate the EOS

We propose the **T-integration method**
to calculate the EOS at fixed scales


T.Umeda et al. (WHOT-QCD), Phys.Rev.D79 (2009) 051501(R)

Our method is based on **the trace anomaly (interaction measure),**


$$\frac{\epsilon - 3p}{T^4} = \left(\frac{N_t^3}{N_s^3} \right) a \frac{d\beta}{da} \left\langle \frac{dS}{d\beta} \right\rangle_{sub}$$

and **the thermodynamic relation.**

$$\frac{\epsilon - 3p}{T^4} = T \frac{\partial(p/T^4)}{\partial T}$$

 $\frac{p}{T^4} = \int_0^T dT' \frac{\epsilon - 3p}{T'^5}$

Test in quenched QCD



[*] G. Boyd et al., NPB469, 419 (1996)

- Our results are roughly consistent with previous results.
- Our results deviate from the fixed $N_t=8$ results [*] at higher T ($aT \sim 0.3$ or higher)
- Trace anomaly is sensitive to spatial volume at lower T (below T_c).
 $V \gtrsim (2\text{fm})^3$ is necessary.

T=0 & T>0 configurations for $N_f=2+1$ QCD

■ T=0 simulation: on $28^3 \times 56$ by CP-PACS/JLQCD *Phys. Rev. D78 (2008) 011502*

- RG-improved Iwasaki glue + NP-improved Wilson quarks



- $\beta=2.05$, $\kappa_{ud}=0.1356$, $\kappa_s=0.1351$

- $V \sim (2 \text{ fm})^3$, $a \approx 0.07 \text{ fm}$, ($m_\pi \sim 634 \text{ MeV}$, $\frac{m_\pi}{m_\rho} = 0.63$, $\frac{m_{\eta_{8S}}}{m_\phi} = 0.74$)

- configurations available on the ILDG/JLDG

■ T>0 simulations: on $32^3 \times N_t$ ($N_t=4, 6, \dots, 14, 16$) lattices

RHMC algorithm, same coupling parameters as T=0 simulation



Formulation for $N_f=2+1$ improved Wilson quarks

$$S = S_g + S_q$$

$$S_g = -\beta \left\{ \sum_{x,\mu>\nu} c_0 W_{\mu\nu}^{1\times 1}(x) + \sum_{x,\mu,\nu} c_1 W_{\mu\nu}^{1\times 2}(x) \right\} \quad \beta = \frac{6}{g^2}$$

$$S_q = \sum_{f=u,d,s} \sum_{x,y} \bar{q}_x^f D_{x,y} q_y^f$$


$$D_{x,y} = \delta_{x,y} - \kappa_f \sum_{\mu} \left\{ (1 - \gamma_{\mu}) U_{x,\mu} \delta_{x+\hat{\mu},y} + (1 + \gamma_{\mu}) U_{x-\hat{\mu},\mu}^{\dagger} \delta_{x-\hat{\mu},y} \right\} - \delta_{x,y} c_{SW} \kappa_f \sum_{\mu>\nu} \sigma_{\mu\nu} F_{\mu\nu}$$

$$c_{SW}(\beta) = 1 + 0.113g^2 + 0.0209g^4 + 0.0049g^6 \quad \text{Phys. Rev. D73, 034501 CP-PACS/JLQCD}$$

$$\frac{\epsilon - 3p}{T^4} = \frac{N_t^3}{N_s^3} \left(a \frac{\partial \beta}{\partial a} \left\langle \frac{\partial S}{\partial \beta} \right\rangle_{sub} + a \frac{\partial \kappa_{ud}}{\partial a} \left\langle \frac{\partial S}{\partial \kappa_{ud}} \right\rangle_{sub} + a \frac{\partial \kappa_s}{\partial a} \left\langle \frac{\partial S}{\partial \kappa_s} \right\rangle_{sub} \right)$$

$$\left\langle \frac{\partial S}{\partial \beta} \right\rangle = N_s^3 N_t \left(- \left\langle \sum_{x,\mu>\nu} c_0 W_{\mu\nu}^{1\times 1}(x) + \sum_{x,\mu,\nu} c_1 W_{\mu\nu}^{1\times 2}(x) \right\rangle + N_f \frac{\partial c_{SW}}{\partial \beta} \kappa_f \left\langle \sum_{x,\mu>\nu} \text{Tr}^{(c,s)} \sigma_{\mu\nu} F_{\mu\nu} (D^{-1})_{x,x} \right\rangle \right)$$

$$\left\langle \frac{\partial S}{\partial \kappa_f} \right\rangle = N_f N_s^3 N_t \left(\left\langle \sum_{x,\mu} \text{Tr}^{(c,s)} \left\{ (1 - \gamma_{\mu}) U_{x,\mu} (D^{-1})_{x+\hat{\mu},x} + (1 + \gamma_{\mu}) U_{x-\hat{\mu},\mu}^{\dagger} (D^{-1})_{x-\hat{\mu},x} \right\} \right\rangle + c_{SW} \left\langle \sum_{x,\mu>\nu} \text{Tr}^{(c,s)} \sigma_{\mu\nu} F_{\mu\nu} (D^{-1})_{x,x} \right\rangle \right)$$

 **Noise method** (#noise = 1 for each color & spin indices)

Beta-functions from CP-PACS/JLQCD results

Trace anomaly needs **Beta-functions** in $N_f=2+1$ QCD

$$\frac{\epsilon - 3p}{T^4} = \frac{N_t^3}{N_s^3} \left(a \frac{\partial \beta}{\partial a} \left\langle \frac{\partial S}{\partial \beta} \right\rangle_{sub} + a \frac{\partial \kappa_{ud}}{\partial a} \left\langle \frac{\partial S}{\partial \kappa_{ud}} \right\rangle_{sub} + a \frac{\partial \kappa_s}{\partial a} \left\langle \frac{\partial S}{\partial \kappa_s} \right\rangle_{sub} \right)$$

Direct fit method Phys. Rev. D64 (2001) 074510

fit $\beta, \kappa_{ud}, \kappa_s$ as functions of $(am_\rho), \left(\frac{m_\pi}{m_\rho}\right), \left(\frac{m_{\eta_{ss}}}{m_\phi}\right)$

$$\begin{pmatrix} \beta \\ \kappa_L \\ \kappa_S \end{pmatrix} = \vec{c}_1 + \vec{c}_2(am_\rho) + \vec{c}_3(am_\rho)^2 + \vec{c}_4\left(\frac{m_\pi}{m_\rho}\right) + \vec{c}_5\left(\frac{m_\pi}{m_\rho}\right)^2 + \vec{c}_6(am_\rho)\left(\frac{m_\pi}{m_\rho}\right) \\ + \vec{c}_7\left(\frac{m_{\eta_{ss}}}{m_\phi}\right) + \vec{c}_8\left(\frac{m_{\eta_{ss}}}{m_\phi}\right)^2 + \vec{c}_9(am_\rho)\left(\frac{m_{\eta_{ss}}}{m_\phi}\right) + \vec{c}_{10}\left(\frac{m_\pi}{m_\rho}\right)\left(\frac{m_{\eta_{ss}}}{m_\phi}\right)$$




$am_\rho \frac{\partial X}{\partial (am_\rho)}$ with fixed $\left(\frac{m_\pi}{m_\rho}\right), \left(\frac{m_{\eta_{ss}}}{m_\phi}\right)$ ($X = \beta, \kappa_{ud}, \kappa_s$)

Beta-functions from CP-PACS/JLQCD results

Meson spectrum by CP-PACS/JLQCD *Phys. Rev. D78 (2008) 011502*.

3 (β) \times 5 (κ_{ud}) \times 2 (κ_s) = 30 data points

fit $\beta, \kappa_{ud}, \kappa_s$ as functions of $(am_\rho), \left(\frac{m_\pi}{m_\rho}\right), \left(\frac{m_{\eta_{ss}}}{m_\phi}\right)$



$$\left(a \frac{\partial \beta}{\partial a}, a \frac{\partial \kappa_{ud}}{\partial a}, a \frac{\partial \kappa_s}{\partial a} \right) \text{simulation point}$$


$$= (-0.334(4), 0.00289(6), 0.00203(5))$$

only statistical error

Trace anomaly in $N_f=2+1$ QCD

$$\frac{\epsilon - 3p}{T^4} = \frac{N_t^3}{N_s^3} \left(a \frac{\partial \beta}{\partial a} \left\langle \frac{\partial S}{\partial \beta} \right\rangle_{sub} + a \frac{\partial \kappa_{ud}}{\partial a} \left\langle \frac{\partial S}{\partial \kappa_{ud}} \right\rangle_{sub} + a \frac{\partial \kappa_s}{\partial a} \left\langle \frac{\partial S}{\partial \kappa_s} \right\rangle_{sub} \right) \quad S = S_g + S_q$$

$$\underbrace{\hspace{10em}}_{(\epsilon - 3p)_\beta/T^4} \quad \underbrace{\hspace{10em}}_{(\epsilon - 3p)_\kappa/T^4}$$




Nt	config. (x 5MD traj.)	
	S_g	$S_q^{(**)}$
56	1300 ^(*)	980
16	1542	647
14	1448	647
12	1492	695
10	863	487
8	628	520
6	657	360
4	802	295


(*) T=0 (Nt=56) by CP-PACS/JLQCD
 S_g calculated with 6500traj.
 (***) thermal. = 1000 traj.

Trace anomaly in $N_f=2+1$ QCD

WHOT-QCD Collaboration




■ peak height = 4~6
in recent Staggered results
($m_q \sim m_q^{\text{phys.}}$)



stout


S. Borsanyi et al.,
arXiv:1007.2580



HISQ

HotQCD,
arXiv:1005.1131

Equation of State in Nf=2+1 QCD



- T-integration

$$\frac{p}{T^4} = \int_0^T dT' \frac{\epsilon - 3p}{T'^5}$$

is performed by the trapezoidal rule (straight line interpolation).

- ϵ/T^4 is calculated from

$$\frac{\epsilon - 3p}{T^4} + \frac{3p}{T^4}$$

- Large error in whole T region

Summary & outlook

We presented the status of WHOT-QCD “EoS” project.

First calculation of EoS with Wilson-type quarks in 2+1 flavor QCD

Small lattice artifacts

- Guaranteed continuum limit
 - No flavor symmetry violations
 - No uncertainty in determination of a LCP
 - $O(a)$ improvement with c_{sw} determined by SF method
 - Simulations on a fine lattice “ $a \approx 0.07\text{fm}$ ”
- } by Wilson-type quarks

Summary & outlook

- Equation of state

More statistics are needed in the lower temperature region

- Results at different scales ($\beta=1.90$ by CP-PACS/JLQCD)
- Techniques to suppress statistical error of EoS

- $N_f=2+1$ QCD just at the physical point

the physical point (pion mass $\sim 140\text{MeV}$) by PACS-CS



$\beta=1.90$ is appropriate to control stat. error at lower T.

Odd Nt config. generation is necessary.


- Finite density

Taylor expansion method to explore EOS at $\mu \neq 0$


Quark mass dependence of Trace anomaly



HotQCD PRD91,054504(2010)



CP-PACS
PRD64,074510 (2001)



HotQCD arXiv1005.1131

QCDFTFD2010

- peak height of the Trace anomaly
→ small quark mass dependence
- Our result seems to be reasonable !
but small m_q is necessary.