

2025年(令和7年)1月4日更新


脊椎動物の脳に存在する新しいエネルギー代謝調節因子(Neurosecretory Protein GL: NPGLと命名)の発見と機能解析

広島大学大学院統合生命科学研究科・総合科学部
教授・Distinguished Professor (DP)
浮穴 和義(うけな かずよし)

研究の要約


- 肥満は糖尿病などの生活習慣病の発症の原因となることから、関連の研究が盛んに進められている。特に、過食やエネルギー収支の破綻により生じることから、摂食中枢が存在する脳内視床下部で作られる摂食関連因子に注目が集まっている。
- 我々は、この視床下部領域に未発見の脳因子があると予測して研究を進めたところ、80アミノ酸残基からなる神経ペプチド(小タンパク質)をコードする遺伝子を鳥類ニワトリの脳で発見し、neurosecretory protein GL(NPGL)と命名した。
- このNPGL遺伝子は脊椎動物に広く保存されていることから、進化的に重要な脳因子であると思われる。そこで、哺乳類のモデル動物であるラットやマウスで機能解析を進めた。
- NPGLの脳室内投与や前駆体遺伝子の過剰発現実験により、NPGLは白色脂肪組織での脂肪蓄積を促すことを見出した。ラットの解析から、食物中の炭水化物から脂肪合成を促すを見出した。一方、マウスの解析から、摂食抑制ニューロンのPOMCニューロンに投射し、摂食行動を亢進したり、エネルギー消費を抑制する可能性がある。この脂肪蓄積作用はニワトリやウズラでも認められることから、脊椎動物に共通した作用であると考えられる。
- さらに、NPGLと進化的に関連があるNPGMというパラログ因子も存在していることを見出しており、2種類の脳因子がどのように協調して作用しているのか、今後の解析が待たれる。
- これまでの解析から、NPGLの脂肪蓄積作用は、悪い肥満ではなく、動物の生命維持に必要不可欠な脂肪蓄積に関係していると考えている。つまり、良い肥満を促す脳因子である。
- 末梢組織での脂肪蓄積を直接制御する脳因子の存在はこれまで知られておらず、神経科学・内分泌学・代謝学の分野で新知見を与える研究になると考えている。
- NPGLに関する研究は、最近巷でも注目されている糖質制限ダイエットや正しい脂肪蓄積の理解へ科学的な知見を与える可能性があると考えている。

肥満やダイエットは、社会的な問題・関心事である



肥満問題の解決には、過食等を引き起こす食欲調節メカニズムの解明が必要

中枢・末梢に発現する生理活性物質が 摂食行動を制御している



この他に機能不明の新しい因子があるのではないか？

学園都市の研究室

第1部 心にプラス

(2)

れない。

こんな事例もあった。ある粉を二ワトリのひなに投与すると食欲が落ち、ネズミに与えると食欲が増進した。機能には謎も多

しみが快感に変わる「ラ

ンナーズハイ」に似た感覚を味わう。脳の不思議に魅せられ、広島大で神経の伝達物質やホルモンの研究を続けてきた。

白い粉の入った容器を

冷蔵庫から得意げに取り

出す。「食欲のコントロ

ールにつながる新しい脳

内物質が絞り込めてき

た」。広島大大学院総合

科学研究科の浮穴准

教授(41)=神経内分泌学

=は容器を指さし、笑み

を浮かべた。

正体はペプチドという

新薬を見据える

基礎研究の先に見据え

るのは、食欲を調節する

新薬の応用研究だ。例え

ば、食欲を増すペプチド

=が分かれれば、ペプチドの

働きを抑えて食欲を減退

させれる薬ができるかもし

らない。こんな事例もあつた。ある粉を二ワトリのひなに投与すると食欲が落ち、ネズミに与えると食欲が増進した。機能には謎も多


いが「薬なら、運動が苦手な人のダイエットに役立つかも」。メタボリック症候群にあらがう現代人を思い、夢を追う。

感情を脳内物質の働きで説明できないか。そんな興味が原点だ。高校時代、剣道の練習中に苦

は、2000年に学会で訪れたインドだった。力レース一歩が1週間続き、「見るのも嫌になつた」。現地の人が毎日食べても平気なのは脳に秘密があるはず、と着想。04年ごろ研究を始めた。

地道な作業はわが身のためでもある。パソコンの前に座る時間が長く、運動不足になりがち。「いつか自分のメタボ対策に役立つ日が来ると信じ、長い間に続けたい」。薬学の専門家や製薬会社との共同研究を思い描き、脳内物質の謎解きに挑む。

食欲の調節



食欲の調節に関連する物質について説明する浮穴准教授



境信重「脱メタボ脳にヒント」
『中国新聞』、2013年6月19日

(境信重)


ニワトリの視床下部漏斗部から、Neurosecretory protein GL (NPGL)と命名した小タンパク質をコードしている前駆体遺伝子を発見した




1. 漏斗部vs小脳でサブトラクション法
2. シーケンス解析(約1000クローニ)
3. シグナルペプチド予測解析
4. ペプチド性因子をコードした遺伝子




BBRC (2014)




既知のペプチド性因子との相同意は全く無い




C末アミド化された神経ペプチド(小タンパク質)
Cys残基が2つあることから内部で架橋



GCE (2017)



NPGL遺伝子は脊椎動物において広く保存されている



アミノ酸配列の分子系統樹解析
(近接結合法)



Gen. Comp. Endocrinol. (2018 review)

Human
Rat
Mouse
Chicken
Western painted turtle
Western clawed frog
Takifugu

Human
Rat
Mouse
Chicken
Western painted turtle
Western clawed frog
Takifugu

Human
Rat
Mouse
Chicken
Western painted turtle
Western clawed frog
Takifugu


Human
Rat
Mouse
Chicken
Western painted turtle
Western clawed frog
Takifugu

MADPGNRRGGIHRPLSFTCSLLIVGMCCVSPFFCHSQT--DLIALSQADPOCWESSSVLLL
MTDPGTRSRNHCPPLSLTCLLIVGMYYVSPVFCHSQT--DLITLNQADPOCWESSSMLLL
MADPGPRSRHSSLIACSLFIVGICYVSPVFCHSQT--DLITLNQADPOCWESSSMLLL
M-DFGNRGRHYNMRILTYSSLVMGVFCVTPSLOCHSQI--DPHALGRADPOCWESSSAVLL
M-DPGNRGRHHS-VRLICSLLIMGVFCVTPFFCHSQN--DPHALGRADPOCWEFSTAALL
M-SSRNKRREYYSMQLTCSLLFLGIFVIPAICHGQV--DPHALGRADPOCWEETSTAILL
M-APSLR-----CLPAAMVLLLGCTCVSSLRDKQVRVDPLSVPQANPHCWKSSSALL

EMWKPRVSNTVSGFWDEIMYLKKSENLKHGALFWDLAQLFWDIYVDCVLSRNHGLGRQL
EMRKPRVSNTVSGFWDEIMYLKKSENLKHGALFWDLAQLFWDIYVDCVLSRNHGLGRREL
EMRKPRVSNTVSGFWDEIMYLKKSENLKHGALFWDLAQLFWDIYVDCVLSRNHGLGRREL
EMRKPRIISDSVSGFWDEIMIFLKKSENLKHGALFWDLAQLFWDIYVDCVLSRNHGLGRQL
EMRKPRIADTVSGFWDEIMYLKKSENLKHGALFWDLAQLFWDIYVDCVLSRNHGLGRQL
EMKPKPRIADTVPAFWDLIMVSLRSSDNKHTALFWDLAQVFWDIYLDCCVLSRNHGLGRQL
EMRVPRIADTVPAFWDLIMVSLRSSDNKHTALFWDLAQVFWDIYLDCCVLSRNHGLGRHV*

VGEEEKLISAAQPQHTRSKQGTYSQLRTSFLKKKELIEDLISMHVRSGSSVIGKVN--L
AGEEQRVSKVLPRHLGIKQGAYSQLRSIYLKKKELIGDLTSMQLQKWGPFGTKEK--L
TGEEEKVSKVLTRHSGIKQGAYSQLLRTLFLKKKELIEDLTSMQMOKRGSRTGKKK--L
AKAQQRITLPSQFTGRNQGMFTHIQRSPVLTKKDFEDLIKNHKHKSRSTLLGRITGEL
AEAEQKITALHSQQLTEQKQGTFSHNQTRVLNKKELIEDLS-IHVHKSGSALLRRVIGGL
NDGNKEITNLISHLTLK--TYSHGQRSILPQKKSLIEELIGIRVHKSESRLLGSIQR--
TALHSLFTDSKSFKLPL-----



EIKRK-- 181
QIKRNRS 183
EVKRK-- 181
GKKRK-- 182
GIKRKQ- 181
GIKRK-- 177
----- 130





ペプチド化学合成により、NPGLを効率よく産出する系を確立した



Microwave-assisted solid-phase peptide synthesizer (Syro Wave, Biotage)



J. Pept. Sci. (2015)





Endocrinology 158:1120-1129 (2017)

Neurosecretory Protein GL, a Hypothalamic Small Secretory Protein, Participates in Energy Homeostasis in Male Mice

Daichi Matsuura,^{1,*} Kenshiro Shikano,^{1,*} Takaya Saito,^{1,*} Eiko Iwakoshi-Ukena,^{1,2} Megumi Furumitsu,¹ Yuta Ochi,¹ Manami Sato,¹ George E. Bentley,² Lance J. Kriegsfeld,³ and Kazuyoshi Ukena^{1,3}

¹Section of Behavioral Sciences, Graduate School of Integrated Arts and Sciences, Hiroshima University, Higashi-Hiroshima 739-8521, Japan; ²Department of Integrative Biology, The Helen Wills Neuroscience Institute, University of California at Berkeley, Berkeley, California 94720; and ³Department of Psychology, The Helen Wills Neuroscience Institute, University of California at Berkeley, Berkeley, California 94720

This report represents the characterization of precursor protein, localization, distribution, and biological action of neurosecretory protein GL (NPGL) in the mouse hypothalamus.



OXFORD
ACADEMIC



ENDOCRINE SOCIETY
JOURNALS

ENDOCRINE
SOCIETY

An Endocrine Society Thematic Issue: Neuroendocrinology 2018

Read our special collection of journal articles published in 2016–2018, focused on neuroendocrinology. Curation of the collection was guided by [Altmetric Attention Scores](#) and Featured Article designations. Clinically relevant topics include a study of patients with pancreatic neuroendocrine tumors, identification of comorbidities associated with acromegaly, analysis of altered HPA axis function in patients with post-traumatic stress disorder, a report of biallelic mutations in KISS1R resulting in normal puberty, and others. Basic science articles highlight the role of PPAR γ in astrocytes, provide a comprehensive review of the genetics of combined pituitary hormone deficiency, describe how a ketogenic diet improves autism-like features, and [report on the distribution of neurosecretory protein GL in the mammalian brain](#). Other basic science contributions describe how oxytocin neurons are activated in the brain, leptin-dopamine signaling in obesity, and more.

【本件リリース先】

文部科学記者会、科学記者会、
広島大学関係報道機関

NEWS RELEASE



広島大学

広島大学広報グループ

〒739-8511 東広島市鏡山 1-3-2

TEL : 082-424-4657 FAX : 082-424-6040

E-mail: koho@office.hiroshima-u.ac.jp

平成 29 年 4 月 28 日

摂食行動やエネルギー代謝調節に関する新しい脳内因子をマウスで発見
-食欲調節に関する脳内基盤の解明に貢献-

【本研究成果のポイント】

- 哺乳類のモデル動物であるマウスの脳から、食欲調節やエネルギー代謝調節に関する脳内因子（Neurosecretory protein GL と命名、略名 NPGL）を世界で初めて発見しました。
- NPGL の発見は、食欲やエネルギー代謝の複雑な調節メカニズムの解明に繋がることが期待されます。
- 今後、我々ヒトの過食や拒食などの摂食調節メカニズムの解明や肥満対策の創薬への応用が期待できます。

2017年4月28日

ScienceDaily

EurekAlert!

AlphaGalileo

ScienceNewslineなど
全世界中で20以上の
サイトから紹介

EurekAlert!
The Global Source for Science News

HOME NEWS MULTIMEDIA MEETINGS PORTALS ABOUT

PUBLIC RELEASE: 28-APR-2017

New appetite control mechanism found in brain

Reason food looks even better when dieting

HIROSHIMA UNIVERSITY



IMAGE: HUNGER AND ENERGY CONSUMPTION MECHANISMS ARE EVEN MORE COMPLEX THAN WE REALIZED. [view more](#)

CREDIT: KAZUYOSHI UKENA

PRINT E-MAIL

RELATED TOPICS

FULL STORY

Health & Medicine

- > Obesity
- > Diet and Weight Loss
- > Nutrition
- > Cholesterol
- Mind & Brain
- > Dieting and Weight Control
- > Nutrition Research
- > Brain Injury
- > Huntington's Disease

Over recent decades, our understanding of hunger has greatly increased, but this new discovery turns things on their head. Up until now, scientists knew that leptin -- a hormone released by fatty tissue, reduces appetite, while ghrelin -- a hormone released by stomach tissue makes us want to eat more. These hormones, in turn, activate a host of neurons in the brain's hypothalamus -- the body's energy control center.

The discovery of NPGL by Professor Kazuyoshi Ukena of Hiroshima University shows that hunger and energy consumption mechanisms are even more complex than we realized -- and that NPGL plays a central role in what were thought to be well-understood processes.

Professor Ukena first discovered NPGL in chickens after noticing that growing birds grew larger irrespective of diet -- suggesting there was more to energy metabolism than meets the eye. Intrigued, the researchers at HU performed a DNA database search to see if mammals might also possess this elusive substance. They found that it exists in all vertebrates -- including humans.

Over recent decades, our understanding of hunger has greatly increased, but this new discovery turns things on their head. Up until now, scientists knew that leptin -- a hormone released by fatty tissue, reduces appetite, while ghrelin -- a hormone released by stomach tissue makes us want to eat more. These hormones, in turn, activate a host of neurons in the brain's hypothalamus -- the body's energy control center.

The discovery of NPGL by Professor Kazuyoshi Ukena of Hiroshima University shows that hunger and energy consumption mechanisms are even more complex than we realized -- and that NPGL plays a central role in what were thought to be well-understood processes.

Professor Ukena first discovered NPGL in chickens after noticing that growing birds grew larger irrespective of diet -- suggesting there was more to energy metabolism than meets the eye. Intrigued, the researchers at HU performed a DNA database search to see if mammals might also possess this elusive substance. They found that it exists in all vertebrates -- including humans.

In order to investigate its role, if any, in mammals, Professor Ukena's team fed three groups of mice, on three distinct diets, to see how NPGL levels are altered. The first set of mice was fed on a low-calorie diet for 24 hours. The second group was fed on a high-fat diet for 5 weeks -- and the third lucky group was fed on a high-fat diet, but for an extended period of 13 weeks.

ScienceDaily*

Your source for the latest research news

Mobile Follow

Breaking News: Learning from

Got Fired from Work? Relax. You may be owed money.

SD Health Tech Enviro Society Quirky

Science News

from research organizations

New appetite control mechanism found in brain

Study explains why food looks even better when dieting

Date: April 28, 2017

Source: Hiroshima University

Summary: A newly discovered molecule increases appetite during fasting, and decreases it during gorging. The neuron-exciting protein, named NPGL -- apparently aims to maintain body mass at a constant, come feast or famine. An evolutionary masterstroke, but not great news for those looking to trim down, or beef up for the summer.

Share:

RELATED TOPICS

FULL STORY

Health & Medicine

- > Obesity
- > Diet and Weight Loss
- > Nutrition
- > Cholesterol
- Mind & Brain
- > Dieting and Weight Control
- > Nutrition Research
- > Brain Injury
- > Huntington's Disease



MULTIMEDIA



Professor Kazuyoshi Ukena, Hiroshima University (IMAGE)



Can You Really Control What You Eat? (IMAGE)

AlphaGalileo

Breaking news for the media. Journalists please register for free access to embargoed news, alerts and contact information.

All regions All categories Science Health Society Humanities Arts Applied science Business

Home

Language English

Content All content

AlphaGalileo

Help

Advertising and Promotion

About us

Prices

Translations

Jobs at AlphaGalileo

Contact us

Conditions of use

News Release



Please register to view contact details

New appetite control mechanism found in brain - reason food looks even better when dieting

28 April 2017 Hiroshima University

A newly discovered molecule increases appetite during fasting - and decreases it during gorging. The neuron-exciting protein, named NPGL - apparently aims to maintain body mass at a constant, come feast or famine. An evolutionary masterstroke, but not great news for those looking to trim down - or beef up for the summer.

Over recent decades, our understanding of hunger has greatly increased, but this new discovery turns things on their head. Up until now, scientists knew that leptin - a hormone released by fatty tissue, reduces appetite, while ghrelin - a hormone released by stomach tissue makes us want to eat more. These hormones, in turn, activate a host of neurons in the brain's hypothalamus - the body's energy control center.

The discovery of NPGL by Professor Kazuyoshi Ukena of Hiroshima University shows that hunger and energy consumption mechanisms are even more complex than we realized - and that NPGL plays a central role in what were thought to be well-understood processes.

Science Newsline Medicine Care > New Appetite Control Mechanism Found... >

Control Mechanism Found in Brain

Published: April 28, 2017.
Released by Hiroshima University



A newly discovered molecule increases appetite during fasting -- and decreases it during gorging. The neuron-exciting protein, named NPGL - apparently aims to maintain body mass at a constant, come feast or famine. An evolutionary masterstroke, but not great news for those looking to trim down -- or beef up for the summer.



Over recent decades, our understanding of hunger has greatly increased, but this new discovery turns things on their head. Up until now, scientists knew that leptin - a hormone released by fatty tissue, reduces appetite, while ghrelin - a hormone released by stomach tissue makes us want to eat more. These hormones, in turn, activate a host of neurons in the brain's hypothalamus - the body's energy control center.

The discovery of NPGL by Professor Kazuyoshi Ukena of Hiroshima University shows that hunger and energy consumption mechanisms are even more complex than we realized. Credit: Kazuyoshi Ukena





Hunger and energy consumption mechanisms are even more complex than we realized. Credit: Kazuyoshi Ukena




マウスにおいて、NPGLは食欲調節中枢において産出される



NPGLは、摂食抑制ニューロンに投射している




Endocrinology (2017)



マウスの脳室内にNPGLを急性投与すると 摂食量が増加した

Endocrinology (2017)





Neurosecretory protein GL stimulates food intake, de novo lipogenesis, and onset of obesity

Eiko Iwakoshi-Ukena^{1,2†}, Kenshiro Shikano^{1†}, Kunihiro Kondo¹, Shusuke Taniuchi¹, Megumi Furumitsu¹, Yuta Ochi¹, Tsutomu Sasaki³, Shiki Okamoto^{4,5,6}, George E Bentley², Lance J Kriegsfeld⁷, Yasuhiko Minokoshi^{4,5}, Kazuyoshi Ukena^{1,7*}

¹Section of Behavioral Sciences, Graduate School of Integrated Arts and Sciences, Hiroshima University, Higashi-Hiroshima, Japan; ²Department of Integrative Biology and The Helen Wills Neuroscience Institute, University of California at Berkeley, Berkeley, United States; ³Laboratory of Metabolic Signal, Institute for Molecular and Cellular Regulation, Gunma University, Maebashi, Japan; ⁴Division of Endocrinology and Metabolism, Department of Homeostatic Regulation, National Institute for Physiological Sciences, Okazaki, Japan; ⁵Department of Physiological Sciences, School of Life Science, Sokendai (The Graduate University for Advanced Studies), Hayama, Japan; ⁶Second Department of Internal Medicine (Endocrinology, Diabetes and Metabolism, Hematology, Rheumatology), Graduate School of Medicine, University of the Ryukyus, Nakagami-gun, Japan; ⁷Department of Psychology and The Helen Wills Neuroscience Institute, University of California at Berkeley, Berkeley, United States

【本件リリース先】

文部科学記者会、科学記者会、
広島大学関係報道機関

NEWS RELEASE



広島大学

広島大学広報グループ

〒739-8511 東広島市鏡山 1-3-2

TEL : 082-424-6762 FAX : 082-424-6040

E-mail: koho@office.hiroshima-u.ac.jp

平成 29 年 8 月 16 日

新規脳内因子が過食と脂肪合成を促し肥満を引き起こすことを発見
-エネルギー代謝調節に関わる脳内基盤の解明に貢献-



【本研究成果のポイント】

- 研究代表者らが発見した脳内因子（Neurosecretory protein GL と命名、略名 NPGL）が哺乳類のモデル動物であるラットにおいて、過食や肥満に関わることを世界で初めて発見しました。
- これまでの研究では、NPGL が食欲調節に関わることまでを明らかにしていましたが、今回の研究では、食欲調節だけでなく、糖質を脂肪に変換し脂肪蓄積を促進するメカニズムに関与していることを解明しました。
- NPGL は炭水化物摂取を促すため、今後、我々ヒトでの糖質制限の科学的理 解や肥満対策の創薬への応用が期待できます。






NPGL及び*NPGL*遺伝子の機能を ラットにおいて解析した

eLife (2017)




Arcuate nucleus


Mediobasal hypothalamus (MBH)




NPGLは食欲調節中枢において産出される



ISH




IHC



Bars = 100 µm



NPGLは、過食を引き起こし、肥満を誘導した





NPGLは過食に加えて、脂肪組織での脂肪合成を亢進させた

eLife (2017)


脂肪合成及び脂肪酸化酵素のmRNA発現量解析



脂肪合成及び脂肪酸化酵素のタンパク質発現解析



脂肪酸不飽和化酵素の活性測定






n=6-7, *P<0.05

NPGLは明期において炭水化物摂取を亢進させた

eLife (2017)

栄養素別摂取カロリー



n = 8, *P<0.05, ***P<0.005





1ケージに3
飼料

飼料の配置は
毎日変更した

NPGLはインスリン低下により発現が誘導された

eLife (2017)



n = 7-8, *P<0.05, **P<0.01


ラットのデータによる考察

eLife (2017)

- 栄養状態が悪い場合:
インスリンの低下が脳に伝わり、
NPGL産生が高まり、摂食行動を
促進させる。
- 栄養状態が良い場合:
食餌由来の脂肪を蓄えるのではなく、新たに生体内で脂肪合成を
亢進させ、脂肪として蓄えさせる
働きがある。



飢餓への対抗・備え




哺乳類のモデル動物であるマウスとラットから得られた成果のまとめと今後の可能性

- 末梢組織での脂肪蓄積を調節するユニークな脳因子の発見である(これまでに報告がない経路)
 - 摂食調節やエネルギー代謝調節の理解へ
- 炭水化物摂取を促す作用を見出しているため、昨今着目されている糖質制限ダイエットへの科学的知見を与えられる
 - 効率の良いダイエット方法を確立できる可能性
- 過度の肥満ではなく、多くの中高年が経験する過体重や隠れ肥満のモデル動物となる可能性がある
 - 内臓脂肪蓄積や異所性脂肪蓄積は代謝疾患の発症を促進させるため、対策が必要
 - NPGLを利用したエネルギー代謝疾患モデル動物を作製し、過食や脂肪蓄積を防ぐ医薬品のスクリーニングへ

哺乳類での結果は鳥類でも同様である

家禽の脳内に存在する神経ペプチドの発見と機能解析

神経ペプチドNPGLは視床下部内で作用し、主に脂肪組織での*de novo*脂肪合成の亢進により、肝臓や脂肪組織での脂肪蓄積を促す新しい脳因子であることを見出している。



我々の発見により、今後下記のような難題が解決できる可能性がある

- ・ 高カロリー食をついつい食べ過ぎてしまうのは何故？
- ・ お腹が空いたときに食べたいと感じるのは何故？
- ・ 食べる量は変わらないのに太ってしまうのは本当？
- ・ 脂vs糖質 どちらが太りやすい？
- ・ 糖質制限はダイエットに有効？ 健康に悪い？
- ・ 糖質を脂肪にする脳内因子や中枢があるの？
- ・ 脂肪が付くのは悪いこと？
- ・ 肥満は防げるの？

SCIENTIFIC REPORTS



OPEN

Localization and function of neurosecretory protein GM, a novel small secretory protein, in the chicken hypothalamus

Received: 8 September 2017

Accepted: 18 December 2017

Published online: 15 January 2018

Kenshiro Shikano¹, Yuki Bessho¹, Masaki Kato¹, Eiko Iwakoshi-Ukena^{1,2}, Shusuke Taniuchi¹, Megumi Furumitsu¹, Tetsuya Tachibana³, George E. Bentley², Lance J. Kriegsfeld⁴ & Kazuyoshi Ukena^{1,4}

Recently, we discovered a novel cDNA encoding the precursor of a small secretory protein, neurosecretory protein GL (NPGL), in the hypothalamic infundibulum of chickens. NPGL plays an important role in the regulation of growth and feeding. A database search indicated that the *NPGL* gene has a paralogous gene: neurosecretory protein GM (*NPGM*), also in chickens. We identified cDNA encoding the *NPGM* precursor in chickens. Morphological analysis showed that *NPGM*-containing cells are specifically localized in the medial mammillary nucleus (MM) and infundibular nucleus (IN) in the hypothalamus. In addition, we found that *NPGM* and *NPGL* are co-localized, especially in the MM. The expression levels of *NPGM* mRNA gradually decreased during post-hatch development, in contrast to those of *NPGL* mRNA. Moreover, we investigated the relationship between *NPGM* and other known factors. *NPGM* was found to be produced in histaminergic neurons in the MM. *NPGM* and histidine decarboxylase, a histamine-producing enzyme, displayed similar expression patterns during post-hatch development. Acute intracerebroventricular injection of *NPGM* decreased food intake, similar to the effect of histamine. To our knowledge, this is the first report of the localization and function of *NPGM* in the brain of vertebrates. These results will further advance the understanding mechanisms underlying energy homeostasis.

【本件リース先】

文部科学記者会、科学記者会、
広島大学関係報道機関

NEWS RELEASE



広島大学

プレスリリース No.3

広島大学広報グループ

〒739-8511 東広島市鏡山 1-3-2

TEL : 082-424-6762 FAX : 082-424-6040

E-mail: koho@office.hiroshima-u.ac.jp

本件の報道解禁につきましては、平成 30 年
1月 15 日(月)19 時以降（日本時間）にお願
いいたします。

平成 30 年 1 月 12 日

記者説明会（1月15日13時・東広島）のご案内

摂食行動を制御する新しい脳内因子を発見 -食欲調節メカニズムの解明や肥満対策の創薬への応用に期待-


【本研究成果のポイント】

- 研究代表者らが最近発見した摂食促進効果のある脳内因子（Neurosecretory protein GL、略名 NPGL）と同じ構造をもった脳内因子 Neurosecretory protein GM（略名 NPGM）が、それとは反対の摂食抑制作用を有することを今回二ワトリの脳で発見しました。
- この NPGM は二ワトリの視床下部領域（※1）に発現しており、中でもヒスタミンニューロン（※2）で産出されることを見出しました。中枢で作られるヒスタミンは、摂食抑制作用や睡眠・覚醒のリズムに関与していることが知られています。
- NPGM もヒスタミンと同様に摂食抑制作用があることを見出しました。今後、ヒトの食欲調節メカニズムの解明や肥満対策の創薬への応用が期待できます。


ニワトリにおいて、NPGLのパラログ因子NPGMを発見し、機能解析を行った

Sci. Rep. (2018)

chicken NPGM	MEFMWKR-R-WYLQ-LGCVLILNLVYA-----NLDYLKEAPASLGOIDHOCWEVSSHGLIVE	53
chicken NPGL	MDFGN-RGRRIHYNMRLTYSL---LVMGVFCVTPSLCHSQIDPLALGRADPOCWESSSAVILLE	58
chicken NPGM	MKKLLKVADPVIALWDFMLFLKESPKPKHNELFNDLAQNFWDMYVDCVLSRSHGMRQLVSP	115
chicken NPGL	MRKPRISDSVSFGEWDFMIFLKSSLENLKHGALFWDLAQLFWDIYVDCVLSRTHGLKGKROLAKA	120
chicken NPGM	RYSSSTYSHRTLQGSAFTNPF	135
chicken NPGL	QQRITTLPSQFTGRNQGMFTHIQRSPVLTKKDFFEDLIKNHKHKSRTLLGRITGELGKKRK	182



NPGM投与後の摂食量



今後の研究展開

- ・ 哺乳類でのNPGMの機能解析
- ・ 少ない飼料で効率よく成長する家禽産出への利用

過食や肥満の治療に期待 食欲抑える物質発見

広島大大学院総合科学研究院科
(東広島市)の浮穴和義教授(46)
〔神経内分泌学〕たちの研究グループがニワトリの脳内に、食欲を抑える物質があることを発見したと15日に発表した。今後、メカニズムの解明に取り組む。研究グループは、ニワトリの視床下部で「NPGM」と呼ばれるタンパク質の一種である物質が分

広島大大学院教授グループ
ニワトリ脳内で分泌

泌されていることを突き止めた。NPGMを人工合成してヒヨコの脳に投与した結果、食欲が減退し摂食量が減った。

浮穴教授たちはこれまでラットの脳内から、食欲を増したり、脂肪の蓄積を促す物質も発見している。

世界的にも注目

早稲田大大学院理工学研究院科の筒井和義教授(65)〔生命理工学〕の話によると、摂食に関する疾患の新たな治療にもつながる可能性があり、世界的にも貢献できる重要な成果である。



浮穴和義教授




鈴中直美「食欲を抑える物質発見」
『中国新聞』、2018年1月16日

Hypothalamic Overexpression of Neurosecretory Protein GL Leads to Obesity in Male C57BL/6J Mice

Yuki Narimatsu^a Eiko Iwakoshi-Ukena^a Keisuke Fukumura^a
 Kenshiro Shikano^{a,b} Megumi Furumitsu^a Masahiro Morishita^a
 George E. Bentley^c Lance J. Kriegsfeld^d Kazuyoshi Ukena^a

^aLaboratory of Neurometabolism, Graduate School of Integrated Sciences for Life, Hiroshima University, Hiroshima, Japan; ^bDepartment of Neurophysiology, Faculty of Medicine, Oita University, Oita, Japan; ^cDepartment of Integrative Biology and the Helen Wills Neuroscience Institute, University of California, Berkeley, CA, USA;
^dDepartment of Psychology, Integrative Biology, and the Helen Wills Neuroscience Institute, University of California, Berkeley, CA, USA



Keywords

Neurosecretory protein GL · Hypothalamus · Obesity · Animal model

Abstract

Introduction: The mechanisms underlying obesity are not fully understood, necessitating the creation of novel animal models for the investigation of metabolic disorders. We have previously found that neurosecretory protein GL (NPGL), a newly identified hypothalamic neuropeptide, is involved in feeding behavior and fat accumulation in rats. However, the impact of NPGL on obesity remains unclear in any animal model. The present investigation sought to elucidate whether NPGL causes obesity in the obesity-prone mouse strain C57BL/6J. **Methods:** We overexpressed the NPGL-precursor gene (*Npgl*) in the hypothalamus using adeno-associated virus in male C57BL/6J mice fed normal chow (NC) or a high-calorie diet (HCD). After 9 weeks of *Npgl* overexpression, we measured adipose tissues, muscle, and several organ masses in addition to food intake and body mass. To assess the ef-

fects of *Npgl* overexpression on peripheral tissues, we analyzed mRNA expression of lipid metabolism-related genes by quantitative RT-PCR. Whole body energy consumption was assessed using an O_2/CO_2 metabolism measurement before an apparent increase in body mass. **Results:** *Npgl* overexpression increased food intake, body mass, adipose tissues and liver masses, and food efficiency under both NC and HCD, resulting in obesity observable within 8 weeks. Furthermore, we observed fat accumulation in adipose tissues and liver. Additionally, mRNA expression of lipid metabolism-related factors was increased in white adipose tissue and the liver after *Npgl* overexpression. *Npgl* overexpression inhibited energy expenditure during a dark period. **Conclusion:** Taken together, the present study suggests that NPGL can act as an obesogenic factor that acts within a short period of time in mice. As a result, this *Npgl* overexpression-induced obesity can be widely applied to study the etiology of obesity from genes to behavior.

【本件リリース先】

文部科学記者会、科学記者会、
広島大学関係報道機関

NEWS RELEASE



広島大学

広島大学広報グループ

〒739-8511 東広島市鏡山 1-3-2

TEL : 082-424-4518 FAX : 082-424-6040

E-mail: koho@office.hiroshima-u.ac.jp

令和3年11月11日

短期間で肥満が誘導されるモデルマウスの作製に成功
—今後の肥満症研究に役立つ成果—

論文掲載

【本研究成果のポイント】

- 近年発見された脳内因子（Neurosecretory Protein GL と命名、略名 NPGL）は食欲調節などのエネルギー代謝調節に関与します。
- NPGL はマウスにおいて短期間で肥満を誘導することを明らかにしました。
- 早期肥満誘導モデル動物を作製できしたことから、今後、ヒトの肥満発症メカニズムの解明や生活習慣病対策への応用が期待できます。

新規脳因子NPGLに関するこれまでの成果 1

1. Kato M, Iwakoshi-Ukena E, Narimatsu Y, Furumitsu M, *Ukena K. Expression of mRNAs encoding hypothalamic small proteins, neurosecretory protein GL and neurosecretory protein GM, in the Japanese quail, *Coturnix japonica*. *Zool. Sci.* 41:50-59 (2024)
2. Narimatsu Y, Kato M, Iwakoshi-Ukena E, Moriwaki S, Ogasawara A, Furumitsu M, *Ukena K. Neurosecretory protein GM-expressing neurons participate in lipid storage and inflammation in newly developed Cre driver male mice. *Biomedicines* 11:3230 (2023)
3. Narimatsu Y, Iwakoshi-Ukena E, Fukumura K, Shikano K, Furumitsu M, Morishita M, Bentley GE, Kriegsfeld LJ, *Ukena K. Hypothalamic overexpression of neurosecretory protein GL leads to obesity in male C57BL/6J mice. *Neuroendocrinology* 112:606-620 (2022)
4. Narimatsu Y, Matsuura D, Iwakoshi-Ukena E, Furumitsu M, *Ukena K. Neurosecretory protein GL promotes normotopic fat accumulation in male ICR mice. *Int. J. Mol. Sci.* 23:6488 (2022)
5. Kato M, Iwakoshi-Ukena E, Narimatsu Y, Furumitsu M, *Ukena K. Effect of stressors on the mRNA expressions of neurosecretory protein GL and neurosecretory protein GM in chicks. *Front. Physiol.* 13:860912 (2022)
6. Naito M, Iwakoshi-Ukena E, Moriwaki S, Narimatsu Y, Kato M, Furumitsu M, Miyamoto Y, Esumi S, *Ukena K. Immunohistochemical analysis of neurotransmitters in neurosecretory protein GL-producing neurons of the mouse hypothalamus. *Biomedicines* 10:454 (2022)
7. Narimatsu Y, Iwakoshi-Ukena E, Naito M, Moriwaki S, Furumitsu M, *Ukena K. Neurosecretory protein GL accelerates liver steatosis in mice fed medium-fat/medium-fructose diet. *Int. J. Mol. Sci.* 23:2071 (2022)
8. Kato M, Iwakoshi-Ukena E, Furumitsu M, *Ukena K. A novel hypothalamic factor, neurosecretory protein GM, causes fat deposition in chicks. *Front. Physiol.* 12:747473 (2021)
9. Fukumura K, Narimatsu Y, Moriwaki S, Iwakoshi-Ukena E, Furumitsu M, *Ukena K. Overexpression of the gene encoding neurosecretory protein GL precursor prevents excessive fat accumulation in the adipose tissue of mice fed a long-term high-fat diet. *Molecules* 26:6006 (2021)
10. Narimatsu Y, Fukumura K, Iwakoshi-Ukena E, Mimura A, Furumitsu M, *Ukena K. Subcutaneous infusion of neurosecretory protein GL promotes fat accumulation in mice. *Helijon* 7:e07502 (2021)
11. Fukumura K, Narimatsu Y, Moriwaki S, Iwakoshi-Ukena E, Furumitsu M, *Ukena K. Effects of overexpression of neurosecretory protein GL-precursor gene on glucose homeostasis and insulin sensitivity in mice. *Int. J. Mol. Sci.* 22:4681 (2021)
12. Fukumura K, Shikano K, Narimatsu Y, Iwakoshi-Ukena E, Furumitsu M, Naito M, *Ukena K. Effects of neurosecretory protein GL on food intake and fat accumulation under different dietary nutrient compositions in rats. *Biosci. Biotechnol. Biochem.* 85:1514-1520 (2021)
13. Kadota A, Iwakoshi-Ukena E, Fukumura K, Shikano K, Narimatsu Y, Furumitsu M, *Ukena K. Effects of irregular feeding on the daily fluctuations in mRNA expression of the neurosecretory protein GL and neurosecretory protein GM genes in the mouse hypothalamus. *Int. J. Mol. Sci.* 22:2109 (2021)

新規脳因子NPGLに関するこれまでの成果 2

14. Shikano K, Iwakoshi-Ukena E, Saito T, Narimatsu Y, Kadota A, Furumitsu M, Bentley GE, Kriegsfeld LJ, *Ukena K. Neurosecretory protein GL induces fat accumulation in mice. *J. Endocrinol.* 244:1-12 (2020)
15. Shikano K, Iwakoshi-Ukena E, Kato M, Furumitsu M, Bentley GE, Kriegsfeld LJ, *Ukena K. Neurosecretory protein GL induces fat accumulation in chicks. *Front. Endocrinol.* 10:392 (2019)
16. Shikano K, Bessho Y, Kato M, Iwakoshi-Ukena E, Taniuchi S, Furumitsu M, Tachibana T, Bentley GE, Kriegsfeld LJ, *Ukena K. Localization and function of neurosecretory protein GM, a novel small secretory protein, in the chicken hypothalamus. *Sci. Rep.* 8:704 (2018)
17. Shikano K, Taniuchi S, Iwakoshi-Ukena E, Furumitsu M, Bentley GE, Kriegsfeld LJ, *Ukena K. Chronic subcutaneous infusion of neurosecretory protein GM increases body mass gain in chicks. *Gen. Comp. Endocrinol.* 265:71-76 (2018)
18. *Ukena K. Avian and murine neurosecretory protein GL participates in the regulation of feeding and energy metabolism. *Gen. Comp. Endocrinol.* 260:164-170 (2018)
19. Shikano K, Kato M, Iwakoshi-Ukena E, Furumitsu M, Matsuura D, Masuda K, Tachibana T, Bentley GE, Kriegsfeld LJ, *Ukena K. Effects of chronic intracerebroventricular infusion of neurosecretory protein GL on body mass and food and water intake in chicks. *Gen. Comp. Endocrinol.* 256:37-42 (2018)
20. Iwakoshi-Ukena E, Shikano K, Kondo K, Taniuchi S, Furumitsu M, Ochi Y, Sasaki T, Okamoto S, Bentley GE, Kriegsfeld LJ, Minokoshi Y, *Ukena K. Neurosecretory protein GL stimulates food intake, de novo lipogenesis, and onset of obesity. *eLife* 6:e28527 (2017)
21. Matsuura D, Shikano K, Saito T, Iwakoshi-Ukena E, Furumitsu M, Ochi Y, Sato M, Bentley GE, Kriegsfeld LJ, *Ukena K. Neurosecretory protein GL, a hypothalamic small secretory protein, participates in energy homeostasis in male mice. *Endocrinology* 158:1120-1129 (2017)
22. Masuda K, Furumitsu M, Ooyama H, Iwakoshi-Ukena E, *Ukena K. Synthesis of neurosecretory protein GM composed of 88 amino acid residues by native chemical ligation. *Tetrahedron Lett.* 57:804-807 (2016)
23. Masuda K, Furumitsu M, Taniuchi S, Iwakoshi-Ukena E, *Ukena K. Production and characterization of neurosecretory protein GM using Escherichia coli and Chinese Hamster Ovary cells. *FEBS Open Bio* 5:844-851 (2015)
24. Masuda K, Ooyama H, Shikano K, Kondo K, Furumitsu M, Iwakoshi-Ukena E, *Ukena K. Microwave-assisted solid-phase peptide synthesis of neurosecretory protein GL composed of 80 amino acid residues. *J. Pept. Sci.* 21:454-460 (2015)
25. *Ukena K, Iwakoshi-Ukena E, Taniuchi S, Bessho Y, Maejima S, Masuda K, Shikano K, Kondo K, Furumitsu M, Tachibana T. Identification of a cDNA encoding a novel small secretory protein, neurosecretory protein GL, in the chicken hypothalamic infundibulum. *Biochem. Biophys. Res. Commun.* 446:298-303 (2014)

学会発表賞受賞 1

1. 12th International Symposium on Avian Endocrinology: Early Career Researcher Runner up Award. Masaki Kato, Eiko Iwakoshi-Ukena, Megumi Furumitsu, Kazuyoshi Ukena “Two Novel Hypothalamic Factors, Neurosecretory Protein GL and Neurosecretory Protein GM, Cause Fat Accumulation in Layer Chicks” イギリス・エдинバラ2022年7月28日
2. 第45回鳥類内分泌研究会 若手研究奨励賞 加藤正暉、岩越栄子、古満芽久美、浮穴和義「ストレス環境下における視床下部分泌性小タンパク質NPGL及びNPGMのmRNA発現変動解析」オンライン2021年12月4日
3. 第45回日本比較内分泌学会大会 学生優秀ポスター発表賞 成松勇樹、加藤正暉、三村朱花、新福夏海、福村圭介、岩越栄子、古満芽久美、森脇翔悟、渡邊大輝、浮穴和義「視床下部分泌性小タンパク質NPGLは骨格筋の量的・質的变化を促し、肥満時の自発運動量を維持する」オンライン2021年11月13日
4. 日本動物学会中国四国支部第72回大会 若手研究者優秀発表賞 成松勇樹、三村朱花、福村圭介、岩越栄子、森脇翔悟、古満芽久美、浮穴和義「視床下部分泌性小タンパク質NPGLが運動量および骨格筋量に与える影響」オンライン2021年6月20日
5. 第22回アディポサイエンスシンポジウム 若手優秀研究奨励賞 鹿野健史朗、松浦大智、齋藤鷹也、越智祐太、古満芽久美、岩越栄子、浮穴和義「視床下部分泌性小タンパク質NPGLはマウスにおいてエネルギー代謝を制御する」大阪2017年8月19日
6. 第34回内分泌代謝学サマーセミナー 優秀ポスター賞 浮穴和義、岩越栄子、鹿野健史朗、近藤邦裕、谷内秀輔、益田恵子、別所裕紀、古満芽久美「視床下部で新たに発見した分泌性小タンパク質の生理機能解析」福岡2016年7月15日
7. 第40回日本比較内分泌学会大会 若手研究者最優秀発表賞 鹿野健史朗、近藤邦裕、益田恵子、岩越栄子、古満芽久美、浮穴和義「Effect of a hypothalamic neurosecretory protein on food preference」広島2015年12月13日
8. 第39回鳥類内分泌研究会 若手研究奨励賞 加藤正暉、鹿野健史朗、益田恵子、岩越栄子、浮穴和義「ニワトリの脳内における2種類の新規視床下部分泌性小タンパク質の生理機能」兵庫2015年11月22日
9. 第46回若手ペプチド夏の勉強会 優秀講演賞 益田恵子、大山晴香、鹿野健史朗、近藤邦裕、古満芽久美、岩越栄子、浮穴和義「マイクロウェーブを用いたFmoc固相法による2つの新規視床下部小タンパク質の合成」京都2014年8月5日
10. 第32回内分泌代謝学サマーセミナー ベストポスター賞 別所裕紀、鹿野健史朗、近藤邦裕、岩越栄子、古満芽久美、谷内秀輔、益田恵子、前嶋 翔、浮穴和義「新規視床下部分泌性小タンパク質の褐色脂肪組織への影響」山梨2014年7月11日
11. 日本動物学会中国四国支部第66回大会 若手研究者優秀発表賞 鹿野健史朗、谷内秀輔、岩越栄子、別所裕紀、前嶋 翔、益田恵子、近藤邦裕、古満芽久美、橘 哲也、浮穴和義「新規視床下部分泌性小タンパク質(NPGM)はニワトリの体重増加を亢進させる」岡山2014年5月11日

学会発表賞受賞 2

12. 第40回日本神経内分泌学会学術集会 若手研究奨励賞 谷内秀輔、鹿野健史朗、近藤邦裕、別所裕紀、前嶋 翔、益田恵子、古満芽久美、岩越栄子、浮穴和義「視床下部で見つけた新規神経ペプチド前駆体遺伝子のラットの成長と脂肪蓄積に対する影響」宮崎2013年10月26日
13. 第36回内藤コンファレンス「分子からみたエネルギーバランスと摂食行動の制御」優秀発表賞 Taniuchi S, Shikano K, Kondo K, Maejima S, Iwakoshi-Ukena E, Ukena K「Functional analysis of a novel hypothalamic neuropeptide gene in the rat」, The 36th Naito Conference on “Molecular Aspects of Energy Balance and Feeding Behavior”, Sapporo, Japan, 札幌2013年9月13日
14. 第28回日本下垂体研究会学術集会 優秀発表賞 谷内秀輔、鹿野健史朗、近藤邦裕、前嶋 翔、益田恵子、別所裕紀、古満芽久美、岩越栄子、浮穴和義「視床下部で発見した新規神経ペプチド前駆体遺伝子の機能解析」岩手2013年8月9日
15. 日本動物学会中国四国支部第65回大会 若手研究者優秀発表賞 別所裕紀、岩越-浮穴栄子、古満芽久美、前嶋 翔、谷内秀輔、橋哲也、浮穴和義「ニワトリで発見した新規神経ペプチドはヒスタミンニューロンに発現している」徳島2013年5月12日
16. 第37回日本比較内分泌学会大会 若手研究者最優秀発表賞 前嶋 翔、岩越栄子、佐藤慧太、坂本浩隆、坂本竜哉、浮穴和義「ラットの視床下部新規神経ペプチドの形態学的解析－既知の生理活性物質との相関と絶食処理に伴う変化－」福井2012年11月30日
17. 日本動物学会中国四国支部第63回大会 若手研究者優秀発表賞 佐藤真実、岩越-浮穴栄子、浮穴和義「マウス視床下部における摂食調節に関わる新規遺伝子のmRNA発現解析」香川2011年5月15日
18. 第35回日本比較内分泌学会大会 若手研究者最優秀発表賞 岩越栄子、田中幸恵、橋哲也、浮穴和義「視床下部における新規摂食調節関連遺伝子の発見」静岡2010年11月20日

特許関係

1. 【取得】国内特許番号P5875105、発明者:浮穴和義・岩越栄子・益田恵子・古満芽久美、発明の名称:ポリペプチド、ポリペプチドの製造方法、摂食調節組成物および摂食量の調節方法、出願人:広島大学、登録日:2016年1月29日
2. 【出願】特願2014-206791、発明者:浮穴和義・岩越栄子・谷内秀輔・鹿野健史朗・近藤邦裕、発明の名称:成長ホルモン発現抑制剤、成人成長ホルモン分泌不全症モデル非ヒト動物、成人成長ホルモン分泌不全症モデル非ヒト動物の作製方法、褐色脂肪細胞の機能調整剤、脂肪蓄積症モデル非ヒト動物及び脂肪蓄積症モデル非ヒト動物の作製方法、出願人:広島大学、出願日:2014年10月7日
3. 【出願】特願2013-219578、発明者:浮穴和義・岩越栄子・谷内秀輔・鹿野健史朗・近藤邦裕、発明の名称:ポリペプチド、核酸、発現ベクター、形質転換体、成長ホルモン発現抑制剤、非ヒト動物及び非ヒト動物の作製方法、出願人:広島大学、出願日:2013年10月22日
4. 【出願】特願2011-248006、特開2012-211121、発明者:浮穴和義・岩越栄子・益田恵子・古満芽久美、発明の名称:ポリペプチド、ポリペプチドの製造方法、摂食調節組成物および摂食量の調節方法、出願人:広島大学、出願日:2011年11月11日、公開日:2012年11月1日

研究費支援

- 科学研究補助金(基盤研究A、基盤研究B、若手研究A、基盤研究C、挑戦的萌芽研究、新学術研究領域、国際共同研究加速基金A、国際共同研究加速基金B)
- 日本学術振興会 特別研究員奨励費
- 国立研究開発法人 農業・食品産業技術総合研究機構 生物系特定産業技術研究支援センター
- 民間研究助成財団(東レ科学振興会、武田科学振興財団、上原記念生命科学財団、三島海雲記念財団、浦上食品・食文化振興財団、小野医学研究財団、鈴木謙三記念医科学応用研究財団、すかいらーくフードサイエンス研究所、花王健康科学研究会、糧食研究会、旗影会、中国電力技術研究財団)

研究室メンバー

現在のメンバー(2024年度)

特任准教授

岩越栄子

特任助教

成松勇樹

研究員

古満芽久美

大学院生

森脇翔悟(D3)

学部生

岡田俊介(B4)



科研費
KAKENHI

研究費(一部)

生研センター

科研費(基盤A, 基盤B, 若手A)

新学術・公募, 国際科研)

民間財団

過去のメンバー

育成助教

福村圭介

ポスドク

谷内秀輔

前嶋翔

研究補助員

大山晴香

田中幸恵

大学院生

鹿野健史朗

加藤正暉

益田恵子

大口悦宏

佐藤真実

近藤邦裕

別所裕紀

松浦大智

越智祐太

齋藤鷹也

門田惇希

内藤万菜

学部生

佐藤瑠奈

山崎玲子

山本拓宜



後藤春菜

三村朱花

新福夏海

中尾ゆきの

小笠原彩乃



Lance Kriegsfeld(UC Berkeley)

George Bentley(UC Berkeley)